

## Part I — Benzene, ethylbenzene and isopropylbenzene

**DOCUMENT DOCUMENT DOCUMENT  
DOCUMENT DOCUMENT DOCUMENT  
DOCUMENT DOCUMENT DOKU  
MENT EΓΓΡΑΦΟ DOCUMENT DO  
UMENTO DOCUMENT DOCUMI  
TO DOKUMENT EΓΓΡΑΦΟ DO  
UMENT DOCUMENTO DOCUME  
T DOCUMENTO DOKUMENTE  
PRAΦΟ DOCUMENT DOCUMEN  
DOCUMENT DOCUMENTO DI  
UMENT EΓΓΡΑΦΟ DOCUMENT  
DOCUMENTO DOCUMENT DOC  
UMENTO DOKUMENT EΓΓΡΑΦΟ**

**This report was requested by the Commission of the European Communities.  
Any views expressed in this report do not necessarily reflect the views of the  
Commission of the European Communities.**

**Cataloguing data can be found at the end of this publication.**

**Luxembourg: Office for Official Publications of the European Communities,  
1993**

**Part I: ISBN 92-826-5894-5  
Parts I-IV: ISBN 92-826-5893-7**


**© ECSC-EEC-EAEC, Brussels • Luxembourg, 1993**

**Reproduction is authorized, except for commercial purposes, provided the  
source is acknowledged.**

***Printed in Belgium***

## Updating of data concerning the impact on the aquatic environment of certain dangerous substances, 2nd part

### Part I Benzene, ethylbenzene and isopropylbenzene



May, 1992

## **PREFACE**

The Commission of the European Communities, Directorate-General for Environment, Nuclear Safety and Civil Protection has commissioned BKH Consulting Engineers to carry out Study Contract B6612-90-006688:

Updating of data concerning the impact on the aquatic environment of certain dangerous substances, 2nd part: benzenes.

The study deals with the following dangerous substances which belong or should belong to List I of Directive 76/464/EEC:

- benzene (7)
- ethylbenzene (79)
- isopropylbenzene (87)
- chlorobenzene (20, 53, 54, 55, 109, 117, 118)
- chloronitrobenzene (27, 28, 29, 30, 63)
- chloroanilines (17, 18, 19, 52)

The study updates information on the description of the substances, the level of contamination of the aquatic environment, the persistence of the substances in the aquatic environment and other media, the toxicity (mainly in the aquatic environment) and bioaccumulation. On the basis of the available data, water quality standards are proposed.

The study is presented in four parts:

Part I	Benzene, ethylbenzene and isopropylbenzene
Part II	Chlorinated benzenes
Part III	Chloronitrobenzenes and 4-chloro-2-nitroaniline
Part IV	Chloroanilines

The study has been carried out by F. Balk, P.C. Okkerman, M. Hof, A. van de Bovekamp and J.W. Dogger. Critical remarks of J. Blok were greatly appreciated.

## SUMMARY

The impact of benzene, ethylbenzene and isopropylbenzene on the aquatic environment is evaluated in order to provide a toxicological basis to derive a proposal for water quality criteria. This evaluation is an update of two earlier reports to EC (Persoone and Vanhaecke, 1982 and Chambon, 1984).

Data on physical and chemical properties, degradability, toxicity, chemical detection and environmental concentrations have been reviewed. The availability of data on the different subjects is summarized in Table i.

Benzene, ethylbenzene and isopropylbenzene are widely used for a variety of products in the chemical industry; they may be used as solvents; benzene and isopropylbenzene are components of gasoline.

Total production in the EC in 1987 of benzene, ethylbenzene and isopropylbenzene is estimated at  $>440 \cdot 10^3$ ,  $2900 \cdot 10^3$  and  $<1370 \cdot 10^3$  tons/year, respectively.

Benzenes can be chemically analyzed by infrared spectroscopy and by gas chromatography. Generally the substances are detected by GC-MS or GC-FID. Detection limits depend on the technique and on the complexity of the sample, but may be less than 1 ng/l.

Recent data on measured concentrations in air, water and soil are lacking. Common concentration of benzene in river water were 0.2 to 3 µg/l. Mean concentrations of ethylbenzene in the river Rhine were 0.3 µg/l with peaks to 2 µg/l. Concentrations in contaminated areas may rise to  $> 5$  mg/kg soil or to  $> 5$  µg/l in groundwater.

As the  $\log K_{ow}$  is low, accumulation on sediment and in animal tissue is low. The bioconcentration factor of benzene for *Daphnia pulex* is 200 and for fish  $<10$ . The chemical reactivity of benzenes in the aquatic environment is low, due to the high stability of the benzene ring system. Benzene and isopropylbenzene are classified as readily biodegradable and also ethylbenzene is completely biodegradable under aerobic conditions. Biodegradation is an important route of elimination from the aquatic system.

Another important route of elimination is volatilization. Calculated half-lives for volatilization from a "standard water system" were less than 6 hours. In air they will degrade rapidly by indirect photolysis with radicals.

Aquatic toxicity data for benzene shows a large variation, with as extremes acute  $LC_{50}$  of 5.3 and 420 mg/l for two fish species. The reproducibility of test results is low: the  $LC_{50}$  in two tests with *Daphnia pulex* differed by a factor 20. The variability in test results may be due, at least partly, to the high volatility of benzene. Information on chronic toxicity is lacking. Studies on early life stages of two marine species showed that embryo development was affected already at 0.05 mg/l.

The acute toxicity of ethylbenzene is comparable to the toxicity of benzene. No chronic toxicity studies were found.

Toxicity data for isopropyl are limited to a mollusc and two crustacean species. The same variability in test results is observed and the toxicity seems to be comparable to the toxicity of ethylbenzene. QSAR estimations of the toxicity based on  $\log K_{ow}$  did not predict the toxicity of ethylbenzene very well, and therefore the QSAR was not considered to be sufficiently reliable to estimate the toxicity of isopropylbenzene.

The  $\text{NOEC}_{\text{ecosystem}}$  was estimated by the modified EPA method: the lowest NOEC for benzene was divided by 10 to arrive at a maximum tolerable concentration of 0.005 mg/l. The toxic effects of benzene, ethylbenzene and isopropylbenzene are expected to be additive. Therefore a water quality criterion of 5  $\mu\text{g/l}$  is proposed for the sum of the three benzenes. This level comes close to the environmental levels cited in the earlier reports as "not uncommon" in river water for benzene and ethylbenzene.

Standards proposed by the USA and Canada for benzene and ethylbenzene range between 2.4 and 5  $\mu\text{g/l}$  and are comparable with the proposed water quality standards (5  $\mu\text{g/l}$ ).

Data on mammalian toxicity are incomplete. A tentative "Tolerable Daily Intake" of 170  $\mu\text{g/kg}$  body weight/day was proposed for benzene and 136  $\mu\text{g/kg}$  body weight/day for ethylbenzene. Benzene is considered to have genotoxic properties; it shows carcinogenic activity in man. For ethylbenzene and isopropylbenzene no data are available.

Based on the proposed water quality standard of 5  $\mu\text{g/l}$  a maximum daily intake by man is estimated at 0.33  $\mu\text{g/kg}$  bodyweight/day, which is far below the above mentioned tolerable daily intake levels.

Table i Information available for benzene, ethylbenzene and isopropylbenzene on the different subjects.

Subject:	BENZENE	ETHYL-BENZENE	ISOPROPYL-BENZENE
1. PHYSICO-CHEMICAL CHARACTERISTICS	+	+	+
Production levels	+	+	+
2. ANALYTICAL DETECTION TECHNIQUES			
Detection methods	+	+	+
Detection levels	+	+	+
3. ENVIRONMENTAL LEVELS			
Residues in the atmosphere	+	+	-
Residues in soil and groundwater	$\frac{+}{-}; \frac{+}{-}$	$\frac{-}{-}; \frac{-}{-}$	$\frac{-}{-}; \frac{-}{-}$
Residues in surface water and sediment	$\frac{+}{-}; \frac{-}{-}$	$\frac{+}{-}; \frac{+}{-}$	$\frac{-}{-}; \frac{-}{-}$
Residues in aquatic organisms	-	+	-
Residues in terrestrial organisms	-	-	-
4. PERSISTENCE AND DEGRADATION PATHWAYS			
Abiotic degradation	+	+	+
Biological degradation and metabolism	+	+	+
5. DISTRIBUTION BETWEEN ENVIRONMENTAL COMPARTMENTS			
Volatization	+	+	+
Sorption	+	+	+
Bioaccumulation:			
Aquatic organisms	+	-	-
Terrestrial organisms	-	-	-
Biomagnification	+	+	+
6. TOXICITY			
Aquatic toxicity:			
Acute toxicity (freshwater; marine)	$\frac{+}{-}; \frac{+}{-}$	$\frac{+}{-}; \frac{+}{-}$	$\frac{+}{-}; \frac{+}{-}$
Chronic toxicity (freshwater; marine)	$\frac{-}{-}; \frac{-}{-}$	$\frac{-}{-}; \frac{-}{-}$	$\frac{-}{-}; \frac{-}{-}$
Quantitative structure activity relations (QSARs)	+	+	+
Toxicity to terrestrial organisms	-	-	-
(semi)field studies	-	-	-
Toxicity to mammals (acute; chronic)	$\frac{+}{-}; \frac{+}{-}$	$\frac{+}{-}; \frac{+}{-}$	$\frac{+}{-}; \frac{+}{-}$
Carcinogenic, mutagenic and teratogenic effects	$\frac{+}{-}; \frac{+}{-}; \frac{-}{-}$	$\frac{-}{-}; \frac{-}{-}; \frac{-}{-}$	$\frac{-}{-}; \frac{-}{-}; \frac{-}{-}$

+ data available  
 $\frac{+}{-}$  insufficient data available  
 - no data available

## ZUSAMMENFASSUNG

Die Wirkung von Benzol, Ethylbenzol und Isopropylbenzol auf die aquatische Umwelt wurde ermittelt mit der Absicht eine toxikologische Grundlage zu beschaffen auf welcher ein Vorschlag für Wasserqualitätskriterien abgeleitet werden könnte.

Dieser Studie stellt eine Aufwertung dar von zwei früheren Berichten an die EG (Persoone und Vanhaecke, 1982 und Chambon, 1984). Werte bezüglich physischen und chemischen Eigenschaften, Abbaubarkeit, Toxizität, chemischer Determinierung und Umweltkonzentrationen wurden untersucht. In Tafel i sind die Ergebnissen dargestellt über die verfügbare Daten der verschiedenen Gegenstände.

Benzol, Ethylbenzol und Isopropylbenzol finden eine breite Anwendung für einen Vielfalt von Produkten der chemischen Industrie; sie können verwendet werden als Lösungsmittel; Benzol, Ethylbenzol und Isopropylbenzol sind Komponenten von Benzin. Die totale Produktion von Benzol, Ethylbenzol und Isopropylbenzol in der EG wird eingeschätzt auf  $>440 \cdot 10^3$ ,  $2900 \cdot 10^3$  und  $1370 \cdot 10^3$  Tonne/Jahr in 1987.

Benzole können chemisch analysiert werden mittels Infrarot-Spektroskopie und durch Gaschromatografie. Im Allgemeinen werden die Stoffen determiniert durch GC-MS oder GC-FID.

Die Determinierungsgrenzwerte sind abhängig von der Anwendungstechnik und von der Komplexizität des Problemusters, können aber weniger als 1 ng/L betragen.

Rezente Werte von gemessenen Konzentrationen in Luft, Wasser und Boden fehlen.

Übliche Konzentrationen von Benzol in Flusswässern betrugen 0,2 bis 3 µg/L. Mittlere Konzentrationswerte von Ethylbenzol im Rhein betrugen 0,3 µg/L mit Höchstwerten bis 2 µg/L. Die Konzentrationen in verunreinigten Gegenden können ansteigen bis  $> 5$  mg/Kg in Böden oder bis  $> 5$  µg/L im Grundwasser.

Akkumulation an Sedimenten und in tierischen Geweben ist niedrig, da die  $\log K_{ow}$  niedrig ist.

Der Biokonzentrationsfaktor von Benzol bezogen auf *Daphnia pulex* ist 200 und für Fische  $< 10$ . Die chemische reaktivierung von Benzole in der aquatischen Umwelt ist niedrig wegen der hohen Stabilität des Benzol-Ringsystems.

Benzol und Isopropylbenzol sind klassifiziert als einfach biologisch Abbaubar und auch Ethylbenzol ist vollständig biologisch abbaubar unter aeroben Verhältnissen.

Biologische Abbaubarkeit ist eine wichtige Beseitigungsweise aus dem aquatischen System.

Eine andere wichtige Beseitigungsweise ist Verflüchtigung. Berechnete Halbwerte durch Verflüchtigung aus ein "standarisiertes Wassersystem" wurden in weniger als sechs Stunden erreicht.

An der Luft werden sie schnellstens abgebaut durch indirekte Photolyse mit Radikalen.

Die Streuung von aquatischen Toxizitätswerten für Benzol variiert bedeutend, mit akuten  $LC_{50}$  zwischen 5,3 und 420 mg/L für zwei Fischarten als äusserste Grenzwerte. Die Reproduzierbarkeit der Versuchsergebnissen ist gering: die  $LC_{50}$  bei zwei Versuchen an *Daphnia pulex* unterschieden sich mit einem Faktor 20.

Die Variationen der Versuchsergebnisse könnten möglicherweise zum Teil zurückzuführen sein auf der hohen Verflüchtigungsgrad von Benzol.



Fakten über chronische Toxizität fehlen. Untersuchungen an frühe Lebensstadien von zwei Marinearten zeigten, daß die embryonale Entwicklung schon beeinflusst wurde bei 0,05 mg/L.

Die akute Toxizität von Ethylbenzol ist vergleichbar mit der Toxizität von Benzol. Es wurden keine Untersuchungsergebnisse über chronischer Toxizität gefunden.

Toxikologische Werte über Isopropyl sind beschränkt auf eine Molluscart und zwei Krustazäenarten. Die gleichen Variationen in den Versuchsergebnissen wurden festgestellt und die Toxizität scheint vergleichbar zu sein mit der Toxizität von Ethylbenzol.

QSAR-Einschätzungen der Toxizität begründet auf  $\log K_{ow}$  ergaben keine deutliche Prognose der Toxizität von Ethylbenzol; deswegen wurde der QSAR nicht als ausreichend zuverlässig betrachtet um die Toxizität von Isopropylbenzol einzuschätzen.

Das  $NOEC_{Ecosystem}$  wurde eingeschätzt durch die angepasste EPA-Methode; die niedrigste  $NOEC$  für Benzol wurde durch 10 geteilt um auf eine maximal akzeptable Konzentration von 0,005 mg/L zu gelangen.

Die toxischen Effekten durch Benzol, Ethylbenzol und Isopropylbenzol sind erwartungsgemäß additiv. Deswegen wird ein Wasserqualitätskriterium von 5 µg/L vorgeschlagen als die Summe der drei Benzole. Dieses Niveau befindet sich nahe an den Umweltniveaus welche in früheren Berichten als "nicht ausserordentlich" in Flusswässern für Benzol und Ethylbenzol genannt wurden.

Das Wasserqualitätskriterium in den USA und in Canada für Benzol und Ethylbenzol variieren zwischen 2,4 und 5 µg/L und sind vergleichbar mit dem vorgeschlagenen Wasserqualitätskriterium (5 µg/L).

Toxizitätswerte über Säugetiere sind unvollständig. Eine vorläufige "akzeptable tägliche Einnahme" von 170 µg/Kg Körpergewicht/Tag wurde vorgeschlagen für Benzol und 136 µg/Kg Körpergewicht/Tag für Ethylbenzol.

Benzol wird betrachtet genotoxische Eigenschaften zu enthalten und deutet auf karzinogenische Aktivitäten an Menschen.

Für Ethylbenzol und Isopropylbenzol sind keine Werten verfügbar.

Ausgehend vom vorgeschlagenen Wasserqualitätskriterium von 5 µg/L wurde die maximale täglich durch Menschen aufzunehmende Menge eingeschätzt auf 0,33 µg/Kg Körpergewicht/Tag, welcher Wert weit niedriger ist als die vorher genannten akzeptablen täglichen Einnahmestufen.

## RÉSUMÉ

L'effet du benzène, de l'éthylbenzène et de l'isopropylbenzène sur l'environnement aquatique a été évalué afin de fournir une base toxicologique d'une proposition concernant les critères de qualité de l'eau. Cette évaluation est un supplément à deux rapports précédents pour la CEE (Persoone en Van Haecke, 1982 et Chambon, 1984). On a rassemblé de l'information sur leurs propriétés physiques et chimiques, la dégradabilité, la toxicité, la détection chimique et les concentrations dans l'environnement. Le tableau i comprend un aperçu de la disponibilité de données concernant ces sujets.

On utilise largement le benzène, l'éthylbenzène et l'isopropylbenzène dans l'industrie chimique. Ils sont utilisés comme dissolvants et le benzène et l'isopropylbenzène sont des composants d'essence.

La production estimée en 1987 dans la CEE de benzène, d'éthylbenzène et d'isopropylbenzène fut respectivement  $> 440 \cdot 10^3$ ,  $2900 \cdot 10^3$  et  $< 1370 \cdot 10^3$  tonnes/an.

Les benzènes peuvent être analysés par spectroscopie infrarouge et par CG. Les méthodes de détection d'usage sont la CG-spectrométrie de masse ou CG-détection à ionisateur de flamme. Les limites de détection dépendent de la technique utilisée et de la composition de l'échantillon, mais elles peuvent être en-dessous de 1 ng/l.

Des données de mesures récentes sur les concentrations dans l'air, dans les eaux de surface et dans les sédiments nous manquent.

Des concentrations habituelles de benzènes dans les eaux fluviales étaient de 0,2 jusqu'à 3 µg/l. Des concentrations moyennes d'éthylbenzènes dans le Rhin étaient de 0,3 µg/l avec pointes jusqu'à 2 µg/l. Dans des régions polluées les concentrations dans la terre peuvent être  $> 5$  mg/kg ou dans les eaux souterraines  $> 5$  µg/l.

Comme  $\log K_{ow}$  est bas, accumulation en sédiments et en tissu animal est minime. Le facteur de bioconcentration de benzènes pour le *Daphnia pulex* est 200 et pour le poisson  $< 10$ . La réactivité chimique de benzènes dans l'environnement aquatique est petite à cause de la grande stabilité du système du cercle benzénique. Le benzène et l'isopropylbenzène ont été classifiés comme aisément biodégradables tandis que l'éthylbenzène est aussi complètement biodégradable dans de conditions aérobies. La dégradation biologique est un procédé d'élimination d'importance dans le système aquatique.

Un autre procédé d'élimination important est la volatilisation. La durée de réduction de moitié qu'on a calculés pour la volatilisation d'un "système aquatique standard" est moins de 6 heures. Dans l'air les combinaisons se dégradent vite par moyen de fotolyse indirecte aux radicaux.

Les données sur la toxicité aquatique des benzènes montrent une variation large avec une valeur ultime de  $CL_{50}$  pour deux espèces de poisson de 5,3 et 420 mg/l respectivement. La reproductibilité des résultats des tests est minime: deux épreuves avec le *Daphnia pulex* donnaient des valeurs de  $CL_{50}$  qui différaient d'un facteur de 20 l'une de l'autre.

Le manque de conformité entre les résultats peut être expliqué, au moins partiellement, par la grande volatilité du benzène.

L'information sur la toxicité chronique fait défaut. Des études de phases de croissance de deux espèces marines ont montré que le développement embryonal fut influencé déjà à un niveau de concentration de 0,05 mg/l.

La toxicité acute d'éthylbenzène est comparable à celle de benzène. Nous n'avons pas trouvé de recherches sur la toxicité chronique.

Des données de la toxicité d'isopropylbenzène sont limitées à celles d'une moule et de deux espèces de crustacés. Elles montrent la même variabilité de résultats des tests et la toxicité semble être comparable à celle d'éthylbenzène.

Des estimations de la toxicité d'éthylbenzène par la Relation Quantitative de la Structure et l'Activité (RQSA), basées sur  $\log K_{ow}$ , ne prédisent pas très bien la réalité; c'est pourquoi le RQSA ne fut pas considéré comme suffisamment fiable de estimer la toxicité d'isopropylbenzène.

$NOEC_{\text{écosystème}}$  a été évalué par la méthode d'EPA modifiée: le NOEC le plus bas pour le benzène fut divisé par dix ce qui eut pour résultat une concentration acceptable au maximum de 0,005 mg/l. On s'attend à ce que les effets de benzène, d'éthylbenzène et d'isopropylbenzène soient additifs. C'est pourquoi on propose une norme de qualité aquatique de 5 µg/l pour l'ensemble des trois benzènes. Cette norme approche celle dont des rapports précédents mentionnent qu'elles ne sont pas singulières pour le benzène et l'éthylbenzène dans les eaux fluviales.

Les normes proposées par les Etats-Unis et le Canada pour le benzène et l'éthylbenzène varient entre 2,4 et 5 µg/l et elles sont comparables à la norme proposée pour la qualité aquatique (5 µg/l).

Les données sur la toxicité pour les mammifères sont incomplètes. Une "prise acceptable par jour" provisoire de 170 µg/kg poids/jour a été proposée pour le benzène et de 136 µg/kg poids/jour pour l'éthylbenzène. On présume que le benzène ait des propriétés génotoxiques; il est cancérogène pour l'homme.

On ne dispose pas de données sur l'éthylbenzène et l'isopropylbenzène.

Si l'on se base sur la norme de qualité aquatique proposée de 5 µg/l la prise maximale par jour pour l'homme sera estimée à 0,33 µg/kg poids/jour ce qui est beaucoup plus bas que les normes citées ci-dessus pour la prise acceptable par jour.

## SAMENVATTING

Het effect van benzeen, ethylbenzeen en isopropylbenzeen op het aquatisch milieu is geëvalueerd om een toxicologische basis te verschaffen aan een voorstel voor waterkwaliteitsnormen. Deze evaluatie vormt een aanvulling op twee eerdere rapporten voor de EEG (Persoone en Vanhaecke, 1982 en Chambon, 1984).

Informatie is verzameld aangaande hun fysische en chemische eigenschappen, afbreekbaarheid, toxiciteit, chemische detectie en concentraties in het milieu. Tabel i bevat een overzicht van de beschikbaarheid van gegevens over deze onderwerpen.

Benzeen, ethylbenzeen en isopropylbenzeen vinden uitgebreide toepassing in de chemische industrie. Ze worden gebruikt als oplosmiddel, en benzeen en isopropylbenzeen zijn bestanddelen van benzine.

De geschatte totale produktie in 1987 in de EG van benzeen, ethylbenzeen en isopropylbenzeen was respectievelijk  $>440 \cdot 10^3$ ,  $2900 \cdot 10^3$ , en  $<1370 \cdot 10^3$  ton/jaar.

Benzenen kunnen chemisch geanalyseerd worden met infraroodspectroscopie en gaschromatografie. Gebruikelijke detectiemethoden zijn GC-MS of GC-FID. Detectiegrenzen hangen af van de gevolgde techniek en van de samenstelling van het monster, maar kunnen lager liggen dan 1 ng/l.

Recente meetgegevens over concentraties in lucht, water en bodem ontbreken. Gangbare concentraties van benzeen in rivierwater waren 0,2 tot 3 µg/l. Gemiddelde concentraties van ethylbenzeen in de Rijn waren 0,3 µg/l met pieken tot 2 µg/l. In verontreinigde gebieden kunnen de concentraties in de grond oplopen tot  $> 5$  mg/kg en in grondwater tot  $> 5$  µg/l.

Accumulatie in sediment en in dierlijk weefsel is gering:  $\log K_{ow}$  is laag. De bioconcentratiefactor van benzeen voor *Daphnia pulex* is 200 en voor vis  $<10$ . De chemische reactiviteit van benzenen in het aquatisch milieu is klein vanwege de grote stabiliteit van het benzeenring systeem. Benzeen en isopropylbenzeen zijn geclassificeerd als goed biologisch afbreekbaar, terwijl onder aerobe condities ook ethylbenzeen volledig biologisch afbreekbaar is. Biologische afbraak is een belangrijk eliminatieproces in het aquatisch systeem.

Een ander belangrijk eliminatieproces is vervluchtiging. Berekende halfwaardetijden voor vervluchtiging uit een "standaard watersysteem" liggen lager dan 6 uur. In de lucht breken de verbindingen snel af via indirecte fotolyse met radicalen.

Aquatische toxiciteits gegevens voor benzenen vertonen grote verschillen, met als uiterste waarden een acute  $LC_{50}$  voor twee vissoorten van respectievelijk 5,3 en 420 mg/l. De reproduceerbaarheid van de testresultaten is laag: twee proeven met *Daphnia pulex* leverden  $LC_{50}$  waarden op die een factor 20 verschilden.

Het gebrek aan overeenkomst tussen testresultaten kan tenminste gedeeltelijk verklaard worden door de grote vluchtigheid van benzeen. Informatie over chronische toxiciteit ontbreekt. Onderzoek aan twee zeewater soorten lieten zien dat de ontwikkeling van het embryo al werd beïnvloed bij concentraties van 0,05 mg/l.

De acute toxiciteit van ethylbenzeen is vergelijkbaar met die van benzeen. Er zijn geen onderzoeken naar chronische toxiciteit gevonden.

Toxiciteitsgegevens voor isopropylbenzeen zijn beperkt tot een mossel en twee kreeftachtigen. Ze laten een zelfde spreiding in testresultaten zien en de toxiciteit lijkt vergelijkbaar met die van ethylbenzeen. QSAR schattingen van de toxiciteit van ethylbenzeen, gebaseerd op de  $\log K_{ow}$ , gaven geen erg goede voorspelling van

de werkelijkheid; de QSAR werd daarom niet voldoende betrouwbaar geacht voor een toxiciteitsschatting van isopropylbenzeen.

De  $\text{NOEC}_{\text{ecosysteem}}$  is geschat met behulp van de gewijzigde EPA methode: de laagste  $\text{NOEC}$  voor benzeen werd door tien gedeeld wat resulteerde in een maximum aanvaardbare concentratie van 0,005 mg/l. De effecten van benzeen, ethylbenzeen en isopropylbenzeen zijn naar verwachting additief. Daarom wordt een waterkwaliteitsnorm voorgesteld van 5  $\mu\text{g/l}$  voor de drie benzenen samen. Deze concentratie benadert die waarvan eerdere rapporten meldden dat ze "niet ongebruikelijk" zijn voor benzeen en ethylbenzeen in rivierwater.

Normen die door de VS en Canada zijn voorgesteld voor benzeen en ethylbenzeen variëren tussen 2,4 en 5  $\mu\text{g/l}$  en zijn vergelijkbaar met de voorgestelde waterkwaliteitsnorm (5  $\mu\text{g/l}$ ).

Gegevens over de toxiciteit bij zoogdieren zijn onvolledig. Een voorlopige "Aanvaardbare Dagelijkse Inname" van 170  $\mu\text{g/kg}$  lichaamsgewicht/dag was voorgesteld voor benzeen en 136  $\mu\text{g/kg}$  lichaamsgewicht/dag voor ethylbenzeen. Van benzeen wordt aangenomen dat het genotoxische eigenschappen heeft; het is bij de mens kankerverwekkend. Voor ethylbenzeen en isopropylbenzeen zijn geen gegevens beschikbaar.

Uitgaande van de voorgestelde waterkwaliteits norm van 5  $\mu\text{g/l}$  wordt de maximum dagelijkse inname door de mens geschat op 0,33  $\mu\text{g/kg}$  lichaamsgewicht/dag. Dit is ver beneden de hierboven genoemde waarden voor de aanvaardbare dagelijkse inname.

## CONTENTS

	Page
PREFACE	i
SUMMARY	ii
ZUSAMMENFASSUNG	v
RESUMÉ	vii
SAMENVATTING	ix
CONTENTS	xi
1. DESCRIPTION	1
1.1 Identification	1
1.2 Physicochemical characteristics	2
1.3 Production levels	2
2. ANALYTICAL DETECTION CHARACTERISTICS	4
2.1 Description of the detection methods	4
2.2 Conclusions	5
3. ENVIRONMENTAL LEVELS	6
3.1 Residues in various environmental compartments	6
3.1.1 Residues in the atmosphere	6
3.1.2 Residues in soil	6
3.1.3 Residues in water	7
3.1.4 Residues in aquatic organisms	7
3.1.5 Residues in terrestrial organisms	7
3.2 Conclusions	8
4. PERSISTENCE AND DEGRADATION PATHWAYS	9
4.1 Abiotic degradation in the aquatic environment	9
4.1.1 Abiotic degradation	9
4.1.2 Conclusions	9
4.2 Biological degradation and metabolism	9
4.2.1 Benzene	10
4.2.2 Ethylbenzene	12
4.2.3 Isopropylbenzene	13
4.2.4 Conclusions	14

CONTENTS (cont'd)

	Page
5. DISTRIBUTION BETWEEN ENVIRONMENTAL COMPARTMENTS	15
5.1 Volatilization	15
5.2 Sorption	16
5.3 Bioaccumulation	17
5.3.1 Aquatic organisms	17
5.3.2 Terrestrial organisms	18
5.4 Environmental distribution	18
6. TOXICITY	19
6.1 Aquatic toxicity	19
6.1.1 Benzene	19
6.1.2 Ethylbenzene	20
6.1.3 Isopropylbenzene	20
6.1.4 Quantitative structure activity relations (QSARs)	28
6.2 Toxicity to mammals	29
6.2.1 Benzene	29
6.2.2 Ethylbenzene	29
6.2.3 Isopropylbenzene	29
6.3 Carcinogenicity, mutagenity and teratogenity	30
6.3.1 Benzene	30
6.3.2 Ethylbenzene and isopropylbenzene	31
7. ENVIRONMENTAL IMPACT ASSESSMENT	32
7.1 Comparison of exposure and effects	32
7.2 Water quality standard	32
7.3 Human exposure	34
REFERENCES	35
APPENDIX 1	41

## 1 DESCRIPTION

Benzene is used as an additive to gasoline and as an intermediate in the synthesis of styrene, synthetic rubber, phenol, cyclohexane, detergents, plastics and resins, nylon and pesticides. It is used as a thinner for lacquers and paints and as a degreasing and cleaning agent. Benzene is also used in dyes, coatings, explosives, flavours and perfumes, photographic chemicals and food processing.

Ethylbenzene is used in the manufacture of styrene and acetophenone, as a solvent in the chemical, paint and rubber industries. It is also used as a constituent of asphalt and naphtha.

Isopropylbenzene is used as a high octane gasoline component; it is used as a thinner for paints and lacquers and it is an important intermediate in phenol manufacture (Sittig, 1981).

### 1.1 Identification

benzene

List I Dir. 76/464/EEC 7


EINECS-no. 2007537

CAS no. 71-43-2

Synonyms: benzeen  
benzol

Molecular formula:  $C_6H_6$

Structural formula:



WLN formula: R

ethylbenzene

List I Dir. 76/464/EEC 79


EINECS-no. 2028494

CAS no. 100-41-4

Synonyms: ethylbenzeen  
ethylbenzol

Molecular formula:  $C_6H_5C_2H_5$

Structural formula:




WLN formula: R2



isopropylbenzene  
List I Dir. 76/464/EEC 87  
EINECS-no. 2027045  
CAS no. 98-82-8  
Synonyms: isopropylbenzeen  
isopropylbenzol

Molecular formula:  $C_6H_5C_3H_7$   
Structural formula:



WLN formula: RY1&1

## 1.2 Physicochemical characteristics

	Benzene	Ethylbenzene	Isopropylbenzene
mol weight	78.12	106.18	120.21
melting point (°C)	5.51	-94.9	-96.0
boiling point (°C)	80.093	136.2	152
density (20°/4°)	0.8794	0.8669	0.864
vapour pressure (mm Hg)	75 (20°C)	10 (25.9°C)	10 mm (38.3°C)
water solubility (20°C) mg/l	1780 <sup>1)</sup>	152	50
volatilization half-life <sup>2)</sup>	5.03	3.1	5.79
" "	4.81		
Log K <sub>ow</sub>	2.19 <sup>3)</sup>	3.15	3.66

1) Keuning and Janssen, 1987.

2) De Bruin, 1985.

3) De Bruijn, 1991.

## 1.3 Production levels

Production levels of benzene, ethylbenzene and isopropylbenzene in EC are presented in Table 1.1., derived from EURECO (1990).

Table 1.1 Production levels of benzene, ethylbenzene and isopropylbenzene in EC countries

Country	Capacity in t/y * actual production						Producer
	Benzene		Ethylbenzene		Isoprop.benz.		
Belgium	130000	y90 [1]					Finaneste NV (Antwerp)
	50000	y90 [1]					SA Sopar NV (Zelzate)
	480000	y90 [1]					BASF (Antwerp)
Denmark							
France	330000	y83 [2,3]	260000 (y87) [2]				CDF Chimie (Carling)
	58000*	y90 [1]					BP Chemical (Laverna)
	150000 (y87) [2]		280000 (y87) [2]				Atochem SA (Gonfreville)
	70000 (y87) [2]						CFR (Gronfreville)
	110000 y83 [3]						Elf Aquitaine (Freyzin)
			6000 (y87) [2]				Elf (Aquitaine Lacq)
			120000 (y87) [2]				Progelec (Le-Pont-de-Claix)
			150000 (y87) [2]				Rhône-Poulenc (Roussillon)
FRG (Germany)	110000	y83 [3]					Deutsche Marathon (Burghausen)
	296000	y90 [1]					VEBA Oel AG (Gelsenkirchen)
	220000	y90 [1]					Redestillationgemeinschaft (Gelsenkirchen)
	200000	y90 [1]					EC Erdoelchemie GmbH (Koeln)
	430000	y90 [1]					Deutsche Shell Chemie GmbH (Godorf)
	195000	y90 [1]					RWE-DEA AG (Wesseling + Heide)
			10000 y90 [1]				RWE-DEA AG (Heide)
			>1000 y90 [1]				BASF AG (Ludwigshafen)
					340000* y90	[1]	Hüls AG (Marl)
						[1]	VEBA Oel (Gelsenkirchen)
Greece							
Ireland							
Italy	750000	y90 [1]					Enmont-Praoil (Prioli+Gela+Porto Torres)
	170000 (y87) [2]						Saras (Sarroch)
	800 y83 [3]						IPP (Shell) (Rho)
			350000 y90 [1]				Enimont Montedipe SpA (Montava)
			40000 (y87) [2]				Selm-Societa E. Montedison (Sarroch)
			110000 (y90) [1]				Enmont-Praoil (Prioli + Sarroch)
					710000 y90	[1]	Enmont-Praoil (Prioli + Porto Torres)
Luxemburg							
Netherlands	815000 (y87) [2,3]		875000 (y87) [2]		24000 (y87) [2]		Dow Chemical BV (Terneuzen)
	285000 y83 [3]						Exxon Chemical Holland (Rotterdam)
	150000 (y87) [3,4]						DSM NV (Geleen)
			375000 (y87) [2]				Shell Nederland BV (Moerdijk)
Portugal	50000 y83 [2]						Petrogal (Porto)
							Companhia Nacional de Petroquímica (Sines)
Spain	80000 y90 [1]						Ertoil (Huelva)
	95000 y90 [1]						CEPSA (Algericas)
	80000 y90 [1]						Repsol Petroleo SA (Puertollano)
	15000 y90 [1]						Ensidesa (Avilas)
			120000 y90 [2]				Repsol Quimica SA (Puertollano)
					170000 y90	[1]	Ertisa (Huelva)
United Kingdom	500000 y90 [1]		32000 y90 [1]		150000 y90	[1]	ICI (River Tees)
	450000 y83 [2,3]						Shell (Blessmere Port)
	145000* y90 [1]						BP (Grangemouth, Wilson)
	170000 y90 [1]						Continental Oil/Conoco (Tamingham)
Total production							
EC	>44000000 (y87) [2]		29000000 (y87) [2]		< 13700000 (y87) [2]		
Total Use EC	64000000 y84 [5]						

1 Information obtained from industry (Eureco, 1990)

2 Water Research Centre (1987)

3 IDET-CEGOS (1983)

4 WHO, 1983

5 Gesellschaft Deutscher Chemiker (1988)

## 2 ANALYTICAL DETECTION CHARACTERISTICS

### 2.1 Description of the detection methods

In the previous evaluation on benzene for the EC, Persoone and Vanhaecke (1982) did not report detection methods for benzene.

In another evaluation on ethylbenzene and isopropylbenzene for the EC, Chambon (1984) reported on the detection methods of ethylbenzene and isopropylbenzene and other aromatic hydrocarbons. Analytical methods are:

- gas chromatography (GC)
- headspace method: detection with photo-ionization detector (PID), flame ionization detector (FID) or mass spectrometry (MS)  
detection limit: 0.1 to 1 µg/l
- direct adsorption: onto a solid phase (e.g. XAD-2 or XAD-7); detection with FID or MS  
detection limit: a few ng to 15 mg/l
- solvent extraction: with e.g. freon, hexane, methylene chloride or pentane  
detection limits: 1 ng/l to 50 µg/l
- infrared spectroscopy (I.R.)
- e.g. extraction with freon  
detection limit: 4 to 40 mg/l

#### Detection in air:

A low level can be detected by filtration of air on charcoal followed by elution with carbon disulphide and detection by GC.

#### Detection in soil:

First, the soil is mixed with freon; after separation the freon is distilled at 70°C. The residue is dissolved in carbon disulphide. Detection by GC or IR.

#### Detection in fish:

A mixed fish sample is vacuum distilled. The volatile substances are trapped on a column which is thermally desorbed. Volatile substances are sent onto a GC capillary column.

#### Detection in water:

The EPA method 602 for purgeable aromates (EPA, 1984) describes a purge-and-trap chromatographic technique for benzene, chlorobenzene, 1,2-dichlorobenzene, 1,3-dichlorobenzene, 1,4-dichlorobenzene, ethylbenzene and toluene in 6 types of water: distilled water, drinking water, surface water, and 3 types of waste water. An inert gas is purged through a contaminated water sample in a chamber with a special temperature. Subsequently the aromates are transferred to the vapour phase and are trapped in a sorbent trap. After purging is completed, the trap is heated and backflushed with the inert gas to desorb the aromates onto a gas chromatographic column. The gas chromatograph is temperature-programmed to separate the aromates which are detected with a photo-ionization detector.

The analytical conditions are:

- temperature-programme: 50°C-2 min., 6°/min to 90°C
- carrier gas: N<sub>2</sub> (20-40 ml/min)
- column packing: 5% SP-1200/1.75% Bentone 34

The detection limits for the EPA method 602 are 0.2 µg/l for both benzene and ethylbenzene with a standard deviations of 10% for benzene and 12% for ethylbenzene.

Geyer (1980) described a detection method for volatile liquid hydrocarbons in seawater, like aldehydes, and aromatic and chlorinated hydrocarbons.

The organics are continuously stripped from a heated, 2-litre sample (70°C) by a pure inert gas (helium). The stripped components are trapped on a Tenax-GC column. Stripping efficiencies were found to be nearly 100%. After stripping, the components trapped on the Tenax-GC, are desorbed by an on-line heating unit (250°C for 15 min.), located outside the GC oven. A pre-column is necessary, since the absorbed compounds are not released simultaneously upon heating. The analysis can be carried out with a GC, provided with the appropriate detector.

The component sensitivity of the entire stripping and analytical method is below 1 ng/l (with a HP 5700A GC with dual FID). The boiling points of the components measured range from about 70-250°C.

## 2.2 Conclusions

The most frequently applied technique for the detection of benzene, ethylbenzene and isopropylbenzene is preceded by a purge and trap technique. As environmental samples often contain a mixture of substances that can disturb the identification of single substances, this pretreatment is important. Generally, the substances are detected by GC-MS or GC-FID.

Detection limits depend on the applied technique and on the complexity of the mixture. In some cases the detection limit can be less than 1 ng/l.

### 3 ENVIRONMENTAL LEVELS

In this chapter a review will be given of the concentration levels for benzene, ethylbenzene and isopropylbenzene in various environmental compartments. These levels depend on local environmental conditions like climate, the organic content of sediments, water flow rate of the rivers, etc. Therefore these levels will give only restricted information of the concentration levels that may occur at other locations.

#### 3.1 Residues in various environmental compartments

##### 3.1.1 Residues in the atmosphere

In the previous evaluation for the EC, Persoone and Vanhaecke (1982) reported benzene concentrations of  $1 \mu\text{g}/\text{m}^3$  to  $50 \mu\text{g}/\text{m}^3$  in air.

In another evaluation for the EC, Chambon (1984) summarized the concentration levels found for ethylbenzene in air. A summary of these levels is presented in Table 3.1.1.

Table 3.1.1 Concentration levels of ethylbenzene in air (Chambon (1984)).

Location	Level ( $\mu\text{g}/\text{m}^3$ )
US urban air	22
tunnel in Amsterdam	13
car park (USA)	480
motorway (USA)	400
The Hague	240
landfill (UK)	3.6 - 49
"	89

No new data of environmental levels in the atmosphere are available.

##### 3.1.2 Residues in soil

Persoone and Vanhaecke (1982) found that data on benzene levels in sediments are scarce and do not permit any interpretation. High concentration in sediments are not expected to occur because of the relatively low sediment/water partition coefficient.

Background concentrations of benzene in soil in The Netherlands varied between  $<0.01$  and  $0.06 \mu\text{g}/\text{kg}$ , but locally concentrations  $>5 \text{ mg}/\text{kg}$  soil were observed on contaminated sites (Slooff, 1988). Measured concentrations in groundwater are usually between  $<0.0005$  and  $0.03 \mu\text{g}/\text{l}$ , but locally concentrations in groundwater may be  $>5 \mu\text{g}/\text{l}$ .

In his evaluation Chambon (1984) reported a concentration of  $0.5 \mu\text{g}/\text{kg}$  (wet weight) for ethylbenzene in sediment from the Los Angeles county.

No new data have been found for concentration level of benzene, ethylbenzene and isopropylbenzene in soil.

### 3.1.3 Residues in water

In their evaluation for the EC, Persoone and Vanhaecke (1982) reported concentration levels of benzene in different types of water. These data are presented in Table 3.1.2.

Table 3.1.2 Concentration of benzene in different types of water (Persoone and Vanhaecke, 1982)

Type of water	Concentration ( $\mu\text{g/l}$ )
drinking water	
- not uncommon	< 0.1 - 1
- maximum	10
river water	
- not uncommon	0.2 - 3
- maximum	100
seawater	
- not uncommon	< 0.05 - 0.2
- maximum	1 - 4
rain water	0.1 - 0.5

In his evaluation for the EC Chambon (1984) reported on the occurrence of ethylbenzene and isopropylbenzene in water. In 1979 ethylbenzene was detected in the river Rhine in levels between 0.7 and 2.3  $\mu\text{g/l}$  (ECETOC, 1988). Chambon (1984) cites a mean level of 0.3 with peaks to 2  $\mu\text{g/l}$  in the Rhine in 1983. Some of the studies on ethylbenzene report that isopropylbenzene was not detected.

Most references report high concentrations in the neighbourhood of gas works. Ethylbenzene is detected in seawater in the neighbourhood of oil platforms (up to 5.5 g/kg). Therefore oil platforms can severely pollute the seawater. On the other hand it was shown that ethylbenzene and isopropylbenzene hardly occur in drinking water.

No new data are available for the levels in water.

### 3.1.4 Residues in aquatic organisms

In his literature study for the EC, Chambon (1984) reported some concentrations in fish from wastewater. Dover sole and croaker contained 4  $\mu\text{g}$  ethylbenzene/kg in the liver. Also in crab traces of ethylbenzene were detected.

No new data are available in aquatic organisms.

### 3.1.5 Residues in terrestrial organisms

No data are available on the concentration levels of benzene, ethylbenzene and isopropylbenzene in terrestrial organisms.

### 3.2 Conclusions

No recent new data have been found on environmental levels of benzene, ethylbenzene and isopropylbenzene in water, sediment and aquatic organisms. The concentrations of ethylbenzene in air are elevated in cities and locations with much traffic. Benzene concentrations in water and soil are usually low but in contaminated areas measured concentrations may rise to  $>5$  mg/kg in soil or to  $>5$   $\mu$ g/l in groundwater.

In general, the lack of data does not allow a sufficient presentation of concentrations of benzene, ethylbenzene and isopropylbenzene in the environment.

## **4 PERSISTENCE AND DEGRADATION PATHWAYS**

### **4.1 Abiotic degradation in the aquatic environment**

#### **4.1.1 Abiotic degradation**

In a previous evaluation for the EC, Persoone and Vanhaecke (1982) reported two references concerning the chemical degradation of benzene in the aquatic environment. They confirmed the minor importance of chemical degradation in the elimination of benzene from the aquatic environment when related to the rapid volatilization of benzene.

In another evaluation for the EC, Chambon (1984) reported that ethylbenzene can be readily oxidized in different solutes by molecular oxygen. Water, however, inhibits this oxidation.

The chemical reactivity of benzene is low, primarily because of the resonance stability of the benzene ring system. As benzene absorbs light at wavelengths below 290 nm, direct photolysis in the watercolumn is not to be expected. Likewise, hydrolysis and oxidation are not expected to be important in natural surface water. In air the benzene derivatives will degrade rapidly by indirect photolysis with radical in the upper layer of the atmosphere (EPA, 1981).

Mill et al. (1980) reported that isopropylbenzene is degraded by alkylperoxyl and hydroxyl free radicals into 2- and 4-isopropylphenol in a 30/70 ratio. In the aquatic environment hydroxyl radicals can be formed from other organic substances, (e.g. riboflavin) and by irradiation with sunlight and therefore degradation along this pathway may occur in water.

No literature was available on hydrolysis of isopropylbenzene in water.

#### **4.1.2 Conclusions**

Limited information is available on the chemical degradation of benzene, ethylbenzene and isopropylbenzene in the aquatic environment. This may be connected to the low reactivity of these substances under environmental aquatic conditions. Furthermore these substances will volatilize rapidly in air because of their high volatilization rate. In air the substances will degrade rapidly by indirect photolysis with radicals.

## **4.2 Biological degradation and metabolization**

The following literature study is based on a review report on biodegradability of priority chemicals by Keuning and Janssen (1987). In an additional search relevant international literature from 1986 until the beginning of 1991 was screened for publications on the biodegradation of benzene, ethylbenzene or isopropylbenzene.

Standardized tests are available to determine biodegradation under aerobic conditions (Annex V in Dir. 79/831/EEC). However, most of the studies reported in literature do not comply with these standardised methods. Therefore, an attempt has been made to classify the substances in line with definitions used by OECD and EC on the basis of the available data on test conditions and results. The following definitions are used:

Primary degradation is the alteration of the chemical structure resulting in the loss of a specific property, or the loss of the parent compound. Ultimate biodegradation is achieved when the test compound is totally utilized by micro-organisms,



producing carbon dioxide, water, mineral salts and new biomass, or mineralization. A substance is classified as readily biodegradable if it shows a positive result in some defined stringent test methods which provide limited opportunities for biodegradation and acclimatization. Readily biodegradable substances are assumed to undergo rapid and ultimate biodegradation in the environment. Substances that biodegrade under more favourable test conditions (prolonged exposure, high biomass concentration, low substance concentration) may be classified as inherently biodegradable. This does not necessarily mean that mineralization is complete, nor that rapid and reliable biodegradation in the environment will occur.

#### 4.2.1 Benzene

##### Biodegradation in standard tests

Benzene was classified as readily biodegradable (OECD TG 301C) by MITI (1985).

##### Biodegradation in the natural environment

Compounds containing an aromatic ring in their structure are wide-spread in nature. Therefore it is not surprising that several native micro-organisms can degrade aromatic compounds like benzene. It is proposed that benzene can be (completely) biodegraded under favourable aerobic conditions in the natural environment (e.g. sufficient nutrients, low concentrations of benzene (< 0,1%) and other (toxic) compounds (Slooff, 1988).

Delfino and Miles (1985) demonstrated that benzene was degraded in groundwater. Under aerobic conditions benzene (1 mg/l) was degraded completely within a period of 16 days. Under anaerobic conditions no degradation was observed within a period of 96 days.

##### Biodegradation under experimental conditions

Several researchers studied the microbial degradation of benzene and isolated micro-organism that can use benzene as the sole source of carbon and energy, i.e. *Pseudomonas putida*, *Moraxella* sp., *Nocardia corallina* (Gibson and Subramanian, 1984).

Oldenhuis et al. (1989) described the effect of addition of *Pseudomonas* strain GJ31 to soil slurries (at 30°C) containing approximately 500 µM benzene.

Without inoculation, the indigenous microflora needed about 30 h for adaptation to degrade chlorobenzene and 100 h for complete removal. Inoculation with cultures of strain GJ31 (5 mg dry weight/l slurry) realized rapid degradation and complete disappearance within 50 h. The results showed that degradation was due to growth of the inoculated cells using benzene as source of carbon and energy. Addition of activated sludge did not stimulate degradation. The degradation rate of aromatic solvents by the added bacteria in soil slurries was similar or higher than that observed in liquid cultures of the same organisms.

Meyer et al. (1984) studied benzene degradation with batch experiments in which sediment of an oil refinery settling pond was used as microbial inoculum. Benzene degraded rapidly when dosed separately, but was not metabolized in the presence of phenol (initial concentrations 100 mg/l). Arvin et al. (1989) found similar results for benzene dosed separately or with other aromatic compounds in a mixed substrate.

Karlson and Frankenberger (1989) assessed the biodegradation of benzene (initial

concentration approximately 450 µg/l) in gasoline contaminated groundwater upon amendment with nutrients ( $\text{NH}_4^-$  /  $\text{NO}_3\text{-N}$ : 100 mg/l) and an enriched mixed cocktail of adapted micro-organisms in batch experiments at 23°C. The benzene content slowly degraded in the unamended sample but dramatically dropped in concentration in the N-amended water and in the inoculated plus N-amended water. In both experiments more than 95% of the benzene in groundwater was removed through microbial action within 73,5 h; a threshold concentration of 22-35 µg/l was observed below which no further degradation took place.

Melcer et al. (1989) measured the variability of influent and effluent concentrations of trace contaminants at a full-scale municipal sewage treatment plant, treating combined domestic and industrial wastewaters. The benzene concentration was reduced from a mean concentration of 0,8 µg/l in the influent by approximately 60% to a mean effluent concentration of 0,3 µg/l.

Aerobic metabolism occurs through the formation of catechol by means of incorporation of molecular oxygen catalyzed by dioxygenases. Subsequently catechols are further degraded by ortho- or meta-cleavage in the endproducts succinic acid and acetyl-CoA (ortho-cleavage) and pyruvic acid and acetaldehyde (meta-cleavage).

Biodegradation of benzene under anaerobic conditions is also possible. Major et al. (1988) investigated the biodegradation of benzene, toluene and xylene isomers (BTX; initial concentration of each compound: 3 mg/l) in anaerobic batch microcosms containing shallow aquifer material at 10°C. After an incubation period of 62 days with nitrate (10 mg/l) the percentage of remaining BTX was 5-15% for the individual compounds compared to the percentage of remaining BTX in the sterile control of 79-86% for the individual compounds. Denitrification was positively confirmed by acetylene blockage of nitrate reductase resulting in accumulation of nitrous oxide.

Hutchins et al. (1991) studied microcosms under anaerobic conditions at 12°C from uncontaminated aquifer material and aquifer material contaminated with JP-4 jet fuel. The microcosms were spiked with nitrate to yield 30 or 75 mg  $\text{NO}_3^-$ /l and several aromatic compounds (final concentration 2-7 mg/l for each compound). In experiments with uncontaminated material benzene was not biodegraded; ethylbenzene biodegraded after a lag period of approximately 30 days to 0.5 mg/l at the end of the 56-day test period. An identical test with contaminated aquifer material showed no removal of benzene and ethylbenzene during the 160 day test period. Wilson et al. (1986) studied the biotransformation of benzene in methanogenic aquifer material. After an incubation period of 40 and 120 weeks (incubation at 17°C in the dark) the benzene initial concentration of 613 µg/l had decreased to 27 and 4% of the sterile control, respectively.

Vogel and Grbic-Galic (1986) observed mineralization of benzene in a mixed culture under anaerobic, methanogenic conditions at 35°C (in the dark). Benzene (3 mM) was used as the sole carbon and energy source. Phenol was the first metabolite detected; the incorporated oxygen atoms originated from water molecules. In another study of Grbic-Galic and Vogel (1987) anaerobic transformation of benzene by mixed methanogenic cultures was described once more. The experiments with this inoculum (10%) were carried out at 35°C in the dark. Benzene was semi-continuously supplied as carbon and energy source in the defined mineral medium. The cultures were fed unlabelled or  $^{14}\text{C}$ -labelled benzene. Gas production was monitored over a period of 60 days. In 4 days the initial

benzene concentration of 15 mM was reduced by 66% to 5 mM with concomitant production of phenol. Complete disappearance of benzene was observed after 34 days. Carbon dioxide and methane were the final products of fermentation. Phenol was found to be the major aromatic intermediate. Additional compounds detected were cyclohexene and propanoic acid.

Schink (1985) carried out methanogenic enrichment experiments using sewage sludge or sediment samples as inoculum (5 ml of anoxic mud per 50 ml enrichment cultures) with hydrocarbons as sole organic substrate. No significant formation of methane was observed in enrichment cultures containing benzene (1 ml/l medium) during 14 weeks of incubation at 28°C and pH 7.2-7.3.

#### 4.2.2 Ethylbenzene

##### Biodegradation in standard tests

No standard tests on the biodegradation of ethylbenzene were reported in the literature.

##### Biodegradation in the natural environment

Almost no data are available in literature on the biodegradation of ethylbenzene in the natural environment. Bouwer (1989) performed aerobic, batch studies at 20°C on samples from a strip-pit pond contaminated with alkylbenzenes and chlorobenzene from tank-truck washing operations. Over a 43-day incubation period  $2 \pm 3\%$  ethylbenzene (initial concentration 100 µg/l) was biotransformed in pondwater. Ethylbenzene was transformed without an apparent lag time in pond water amended with nutrients (10 mg acetate/l; 5 mg N/l; 5 mg P/l) as concentrations were below the detection limit after 12 days of incubation. The addition of nutrients and sediment collected from the bottom of the strip-pit pond (concentration 0.61 g/l) to the pond water produced similar rapid biotransformation. In a repeated batch experiment with pondwater (with and without sediment) amended with nutrients, ethylbenzene (100 µg/l) was completely biotransformed within 24 hours.

In an experimental aerobic activated sludge installation with an influent concentration of 60 mg/l, Kincannon et al. (1983) observed a removal percentage of more than 99% for ethylbenzene. Evaporation and adsorption to activated sludge were responsible for a removal of 5.2-7.5% and 0-0.19%, respectively. Apparently the remaining 92.4-94.6% was removed by microbial activity.

##### Biodegradation under experimental conditions

Biodegradation of ethylbenzene under aerobic conditions by several micro-organisms is described. Gibson (1972) already described the metabolism via 3-ethylcatechol by *Pseudomonas* sp. and a *Nocardia* sp. In other studies by Gibson et al. (1973, 1984) the metabolism of ethylbenzene (100 mg/l) by *Pseudomonas putida* was investigated. The major metabolic route took place via 1,2-dihydroxy-3-ethylcyclohexa-3,5-diene and cis-1,2-dihydroxy-3-(1-hydroxyethyl)cyclohexa-3,5-diene followed by ring fission and further metabolism.

The minor route was via the metabolite 1-phenylethanol which was degraded to the corresponding catechol and subsequently transformed via meta ring-fission.

Smith and Ratledge (1989) reported on catabolism of alkylbenzenes by *Pseudomonas* sp. NCIB 10643. *Pseudomonas* sp. NCIB 10643 grew on a range of mono-substituted alkylbenzenes (C<sub>2</sub>-C<sub>7</sub>), including ethylbenzene and isopropyl-

benzene. Both alkylbenzenes were metabolized via ring attack, proceeding via initial dioxygenase activity resulting in 3-alkyl-substituted catechols subsequently followed by the meta ring-cleavage pathway.

Utkin et al. (1991) isolated *Pseudomonas* sp. Y2 from oil-contaminated soil in the presence of ethylbenzene as a sole carbon and energy source. Strain Y2 transformed ethylbenzene via the metabolites 2-phenylethanol and phenylacetate and also via 1-phenylethanol. Detection of catechol-2,3-dioxygenase in cell-free extracts of strain Y2 indicated that ethylbenzene could be degraded via the corresponding catechols and subsequently the meta ring-cleavage pathway.

Holland et al. (1985) described the metabolism of ethylbenzene (0.5-1 mg/l) by the fungus *Mortierella isabellina*. As metabolite phenylethanol was detected.

Bouwer and McCarty (1985) studied aerobic columns with immobilised micro-organisms growing on 1 mg acetate/l, which degraded low ethylbenzene concentrations (influent concentration 9.1 µg/l). When the columns were operated with a retention time of 20 minutes the removal of ethylbenzene was 99%.

Few data are available on anaerobic degradation of ethylbenzene. Wilson et al. (1986) studied the biotransformation of ethylbenzene in methanogenic aquifer material. After an incubation period of 20 weeks, 20°C in the dark, no degradation of ethylbenzene (269 µg/l) was observed; after 40 weeks of incubation the ethylbenzene concentration had decreased to 69% of the sterile control.

Bouwer and McCarty (1983) observed no transformation of ethylbenzene (40-114 µg/l) under anaerobic denitrifying conditions with activated sludge as inoculum at 25°C during an incubation period of 11 weeks.

Bouwer (1985) described that immobilised micro-organisms in anaerobic methanogenic columns, growing on acetate as the primary substrate could not degrade simultaneously added ethylbenzene (initial concentration 12 µg/l).

#### **4.2.3 Isopropylbenzene**

##### **Biodegradation in standard tests**

Isopropylbenzene was classified as readily biodegradable (OECD TG 301C) by MITI (1985).

##### **Biodegradation in the natural environment**

No data are available in literature on biodegradation of isopropylbenzene in the natural environment. However, as substances containing an aromatic ring structure are wide-spread in nature, it is assumed that isopropylbenzene can be biodegraded under aerobic conditions by a large variety of micro-organisms.

##### **Biodegradation under experimental conditions**

Information on the biodegradation of isopropylbenzene under experimental conditions is limited. Based on the simple aromatic structure it is generally assumed that isopropylbenzene is biodegraded under aerobic conditions, but that the isopropyl moiety is hindering microbial oxidation resulting in lower degradation rate. Isopropylbenzene for example was degraded a factor 2 slower than toluene by a toluene-grown culture of *Pseudomonas putida* (Gibson, 1972; Gibson and Subramanian, 1984). Proposed is that degradation occurs via 3-alkyl-substituted catechols.

Smith and Ratledge (1989) reported on catabolism of alkylbenzenes by *Pseudomonas* sp. NCIB 10643. *Pseudomonas* sp. NCIB 10643 grew on a range of mono-substituted alkylbenzenes ( $C_2$ - $C_7$ ), including ethylbenzene and isopropylbenzene. Both alkylbenzenes were metabolized via ring attack, proceeding via initial dioxygenase activity resulting in 3-alkyl-substituted catechols subsequently followed by the meta ring-cleavage pathway.

No data are available in literature on the biodegradation of isopropylbenzene under anaerobic conditions.

#### 4.2.4 Conclusions

As substances containing an aromatic ring structure are wide-spread in nature, it is assumed that substances with a similar structure like benzene, ethylbenzene and isopropylbenzene, are biodegradable by a variety of micro-organisms.

Benzene is classified as readily biodegradable. Anaerobic biodegradation was reported but at a very low rate and after a long lag phase.

Under aerobic conditions ethylbenzene is completely biodegradable. Biodegradation under anaerobic conditions does not seem to be important.

Isopropylbenzene is classified as readily biodegradable. Biodegradation under anaerobic conditions has not been reported.

## 5 DISTRIBUTION BETWEEN ENVIRONMENTAL COMPARTMENTS

### 5.1 Volatilization

In a previous evaluation on benzene for the EC, Persoone and Vanhaecke (1982) found an evaporation half-life of 37 minutes in a water column of 1 meter depth at 25°C. They also reported on the rapid volatilization of benzene from oil spills.

In another evaluation for the EC, Chambon (1984) found that ethylbenzene evaporates from water with a half-life of 7.5 hours for a water depth of 1 meter at 25°C. Under the same conditions a half-life of 5.79 hours was reported for isopropylbenzene.

According to Mackay and Leikonen (1975) an evaporation half-life for benzene was 4.81 h at 25°C and 5.03 h at 10°C.

Hellman (1987) studied the vaporization behaviour of benzene. He developed a model for the quantification of the volatilization rate in the environment. With this model experiment a volatilization rate of  $11.6 \cdot 10^{-3} \text{ min}^{-1}$  was found.

As benzene volatilizes rapidly most degradation takes place in air where it is degraded quickly:  $t_{1/2} < 1 \text{ day}$  (Ware 1980).

Ince and Inel (1989) carried out laboratory experiments to measure the volatilization rate of different substances. For ethylbenzene a volatilization (first-order) rate constant of  $0.71 \text{ h}^{-1}$  (25°C) was found. Aeration of the solution resulted in a doubling of the volatilization rate.

No literature was available on the volatilization of isopropylbenzene from water.

In recent years several models have been developed to estimate the volatilization behaviour of substances in the natural environment. In this evaluation study the estimation method developed by De Bruin (1985) is used to estimate the volatilization half-life ("standard half-life"). The calculation is based on physical parameters like water solubility, vapour pressure, water speed (1 m/s), air velocity (3 m/s) and water depth (1 m). The calculation method is presented in Appendix 1. The calculated half-lives for the benzene, ethylbenzene and isopropylbenzene are presented in Table 5.1.1.

Table 5.1.1 Calculated volatilization rate (standard half-life) for benzene, ethylbenzene and isopropylbenzene, according to De Bruin (1985).

Substance	Half-life (h)
benzene	5.0
ethylbenzene	3.1
isopropylbenzene	5.8

A summary of published data on volatilization rates and half-lives data is given in Table 5.1.2.

Table 5.1.2 Volatilization rates and half-lives reported for benzene, ethylbenzene and isopropylbenzene.

Substance	Volatilization rate (h <sup>-1</sup> )	Half-life (h)	Ref.
benzene	1.1	0.62	1
	4.8	0.14	2
	0.70	1.0	3
ethylbenzene	0.092	7.5	4
	0.71	0.98	5
isopropylbenzene	0.12	5.79	4

References:

- 1 Persoone and Vanhaecke (1982)
- 2 Mackay and Leikonen (1975)
- 3 Hellmann (1987)
- 4 Chambon (1984)
- 5 Ince and Inel (1989)

Table 5.1.2 shows a high variety in the various volatilization half-lives of each substance. This is mainly caused by the difference in experimental conditions. Differences in half-lives of the various compounds will be caused by the specific physical and chemical properties of the substances

For a proper comparison of the volatility of the substances, Table 5.1.1 should be used, because those calculations are based on identical physical and chemical properties and assumptions.

In general, it can be concluded that the substances will volatilize rapidly from water into the air.

## 5.2

### Sorption

For the fate of a substance in the environment, sorption can play an important role. Substances may adsorb on suspended particles and on sediment where they can remain for a long time. After adsorption a substance may return to the aquatic environment by desorption. It is possible to measure a distribution coefficient of a substance between sediment and water. This parameter is called the sediment/water partition coefficient. The value of this coefficient is dependant on the physical and chemical properties of the substance, e.g. solubility and on the organic carbon content of the sediment. Usually, the partition coefficient is calculated for the carbon content of the sediment. This parameter is called the "organic carbon partition coefficient" ( $K_{oc}$ ).

For many substances the  $K_{oc}$  has been measured. For other substance the  $K_{oc}$  can be estimated from the octanol/water partition coefficient ( $K_{ow}$ ) or from the water solubility.

For benzene, ethylbenzene and isopropylbenzene these partition coefficients and the solubility in water are presented in Table 5.2.1 (data have been derived from chapter 1)

Table 5.2.1 Physical properties of benzene, ethylbenzene and isopropylbenzene

Parameter	Benzene	Ethylbenzene	Isopropylbenzene
solubility (mg/l)	1800	152	50
H (atm·m <sup>3</sup> /mol)	5.4·10 <sup>-3</sup>	5.8·10 <sup>-3</sup>	1.28·10 <sup>-3</sup>
log K <sub>oc</sub>	1.87	1.98	
log BCF	1.34		
vol. half-life (h <sup>-1</sup> )	5.0	3.1	5.8

As a result of the low K<sub>ow</sub>-value and low K<sub>oc</sub>-value it is not to be expected that adsorption will play a major role in the distribution in the environment.

### 5.3 Bioaccumulation

#### 5.3.1 Aquatic organisms

If experimental data are absent, the potential of an organic substance to bioaccumulate may be estimated from its water-octanol partition coefficient, expressed as log K<sub>ow</sub>, see chapter 1.2. If log K<sub>ow</sub> is above 3, the substance has the potential to accumulate in the biomass. Whether accumulation actually occurs, depends among other things on the capacity of the organisms to metabolise or eliminate the substance. The log K<sub>ow</sub> for benzene is 2.2, for ethylbenzene 3.2 and for isopropylbenzene 3.7.

The data collected by Persoone and Vanhaecke (1982) indicate that the accumulation level of benzene in fish is restricted. Extreme values were 0.2 and 135, but most values ranged between 1 and 10. Data for other species were not available at the time.

Only few new data were found for benzene, see Table 5.3.1. All reported bioconcentration factors (BCF) were based on wet weight.

Two BCFs were found for algae: 30 for *Chlorella fusca* var. *vacuolata* and 2217 for *Selenastrum capricornutum*. Exposure of *Daphnia pulex* to two different concentrations of benzene (40 and 80 µg/l) resulted in BCFs of 225 and 203, respectively. For the fish *Leuciscus idus melanotus* exposure to benzene resulted in a BCF less than 10. Apparently, benzene accumulates (or adsorbs) in algae and in *Daphnia* to a higher level than in the tested fish species. The data confirm the earlier statement in Persoone and Vanhaecke (1982) that bioaccumulation of benzene in fish is not important. Probably the fish is capable to eliminate or metabolize benzene.

For ethylbenzene and isopropylbenzene no data on bioaccumulation were available from the study of Chambon (1984), nor could new data be added in the present update. However, on the basis of the similarity of these substances in chemical characteristics, the similarity in biodegradability and in toxicity (see Chapter 6) it is expected that bioaccumulation in fish will not be important.



Table 5.3.1 Bioconcentrationfactors for benzene

Substance/ species	Lifestage age/size	Conc. (µg/l)	Test system	Uptake period	BCF (wet weight)	Reference
Algae						
<i>Chlorella fusca</i> <i>var. vacuolata</i>		50	S	24 h	30	Freitag, 1985
<i>Selenastrum</i> <i>capricornutum</i>		10000	S	24 h	2217	Casserly, 1983
Crustaceans						
<i>Daphnia pulex</i>	1.9-2.1 mm	40	S	24 h	225	Trucco et al., 1983
		80	S	24 h	203	
Fish						
<i>Leuciscus idus</i> <i>melanotus</i>		50	S	24 h	< 10	Freitag, 1980

### 5.3.2 Terrestrial organisms

No references were found on bioaccumulation of benzene, ethylbenzene and isopropylbenzene in terrestrial species.

### 5.4 Environmental distribution

The fate of a substance in the aquatic environment is dependant on a number of physical and chemical characteristics such as volatilization, sorption, bioaccumulation, chemical and biological degradation.

Volatilization, sorption and bioaccumulation have been discussed in this chapter, and chemical and biological degradation was discussed in chapter 4.

After release into the environment benzene, ethylbenzene and isopropylbenzene will mainly partition to the atmosphere because of their high volatilization rate. The bioaccumulation of benzene in fish is not important. Few data are available on adsorption and bioaccumulation of ethylbenzene and isopropylbenzene. The chemical degradation rate in water will be low as a result of the stability of these aromates, but the substances are completely biodegradable in water under aerobic conditions.

Sorption, bioaccumulation and biodegradation, however will be of minor importance because of the short life time in the aquatic environment.

## 6 TOXICITY

A general remark should be made with regard to the interpretation of the data presented in the literature. Benzene, ethylbenzene and isopropylbenzene are highly volatile substances, indicated by half-lives less than 6 hours (Table 5.1.1). As a consequence these substances will also disappear from the test vessels in toxicity studies. According to Persoone and Vanhaecke (1982) the concentration decrease in static tests is around 25% after 1 day and may rise to 80 - >99% after 4 days, depending on the shape and surface to volume ratio of the container, agitation of the system, the use of a cover and the type of the test (static or flow-through).

As the present report is an update of two EC-reports (Persoone and Vanhaecke, 1982; Chambon, 1984), a search was carried out for toxicity data published since 1980. Since then only a limited number of studies have been reported. Also in most of these new studies, test were static and test substance concentrations have only been reported as nominal concentrations, and should therefore be interpreted with caution. The data are presented in tables 6.1.1 to 6.1.3. A selection of data from the previous EC-reports is included in the tables.

### 6.1 Aquatic toxicity

#### 6.1.1 Benzene

According to Persoone and Vanhaecke (1982) the toxicity of benzene for micro-organisms varied between 50 and 200 mg/l. However, benzene may interfere with the chemotaxis of micro-organisms at relatively low concentrations, e.g. 0.1 mg/l. However, the chemotaxis inhibition is a reversible process.

No new data were found on the toxicity of benzene to micro-organisms.

The data for algae as reported by Persoone and Vanhaecke (1982) vary largely. In their evaluation they state that maximum no effect levels for marine algae species vary between 10 and 50 mg/l. Compared to this range, the new data for *Ankistrodesmus falcatus* and *Selenastrum capricornutum* seem to be very high. The MATC of < 34 mg/l for the red algae *Champia parvula* (14 days renewal test) falls well within the above mentioned range.

The mean LC<sub>50</sub>-value for freshwater invertebrates calculated by Persoone and Vanhaecke (1982) approximated 260 mg/l, with the rotatorian as the most sensitive species (14.4 mg/l). Marine crustacean species (mean LC<sub>50</sub> 67 mg/l) were more sensitive to benzene than the mollusc.

New data were reported for 9 freshwater vertebrates, including two insects, three protozoans, three hydrozoans and a snail. These seem to fit well in this range.

However, also in these data a large intra-species variability in the results is observed (e.g. a factor 20 for the toxicity of *Daphnia pulex*).

For one species extremely sensitive life stages are reported: in tests with the marine echinoderm *Paracentrotus lividus* both the sperm and the embryo were affected at 0.078 mg/l. Endpoints were developmental effects (gut and skeletal malformations), fertilization success and chromosomal abnormalities (genotoxic effect). This low value compares well with the value cited by Persoone and Vanhaecke (1982) for the lowest concentration (0.053 mg/l) for "slight" effect in the embryos of a marine fish.

For most fish species the LC<sub>50</sub> fell within the range of 5 to 50 mg/l (Persoone and Vanhaecke, 1982). Almost no change in LC<sub>50</sub>-values was found between 24 h and 96 h tests. The new data found in this study for *Oryzias latipes* and *Poecilia reticulata*

seem to be rather high as compared to this range. The sensitivity of two amphibian species tested by the same author were of the same order of magnitude. It is not clear whether benzene concentrations were maintained during these studies.

The new data on benzene include data on a number of new species of freshwater invertebrates and amphibians. The toxicity values fall well within the range of data in the previous report. The lack of studies on chronic toxicity as observed in 1982 is confirmed in the present report. Two studies on early life stages (in an echinoderm and a fish species) showed that the embryo development was affected at concentrations around 0.05 mg/l.

#### **6.1.2 Ethylbenzene**

In the previous study on ethylbenzene for EC Chambon (1984) reported toxicity data for bacteria, algae, invertebrates and fish.

The data compare well to the data for benzene, and also the same wide variability is observed which may, at least partly, be connected to the high volatility of the test substance.

Only for two species new toxicity data were reported since 1980. The  $LC_{50}$ -values for the crustaceans were 2.1 mg/l for *Daphnia magna* being 40 to 50 times lower than the older values, and 15.4 mg/l for *Artemia*. The latter value fits well into the range of earlier data for marine crustaceans.

Chronic studies with ethylbenzene have not been found.

#### **6.1.3 Isopropylbenzene**

In the previous study on isopropylbenzene for EC (Chambon, 1984) toxicity data were available for *Daphnia magna* and *Mytilus edulis*. In addition new data were found on *Daphnia magna* and *Artemia* (Table 6.3.1). The newer data for *D. magna* are more than 2 orders of magnitude lower than the older (0.6 versus 95 mg/l). It should be remarked that the latter concentration (95 mg/l) is higher than water solubility of isopropylbenzene in water: 50 mg/l.

Data on chronic toxicity are limited to an unspecified test with the mussel larvae.

The limited number of data available suggest that the acute toxicity is comparable to the toxicity of ethylbenzene. Chronic data are lacking.

Table 6.1.1 Single species toxicity data for benzene

Species	Lifestage age/size	A/N	Test system	Purity/ solvent	Testwater	pH	Hardness mg CaCO <sub>3</sub> /l	Expo time (h)	Parameter	Results (mg/l)	Reference	Quality
Algae Ankistrodesmus falcatus		N	S	Pure	CHU 10 medium	8		24	EC <sub>50</sub> growth	310	Wong, 1984	2-3
Champia parvula (M)		N	R		Sea water + suppl.			14d	MATC growth, reproduction	<34.2	Thursby et al., 1985	
Selenastrum capricornutum		N	S					96	NOEC growth	600	Stooff, 1983b	
Protozoans Tetrahymena ellioti			S					24	NOEC survival	391	Rogerson et al., 1983	
Crustaceans Artemia	Larvae		S	Purity >97%				24	LC <sub>50</sub>	127.3	Abernethy et al., 1986#2	
Daphnia cucullata	≤ 24 h	N	S					48	LC <sub>50</sub>	373	Stooff, 1983b	
Daphnia magna			S	Purity		6-7		48	LC <sub>50</sub>	31.2	Bobra, 1983	
Daphnia magna	Neonates		S	Purity >97%				48	LC <sub>50</sub>	127.3	Abernethy et al., 1986#2	
Daphnia magna	≤ 24 h	N	S					48	NOEC	240	Stooff, 1983b	
Daphnia magna	≤ 24 h	N	S	>97%				48	LC <sub>50</sub>	400	Stooff, 1983b	
Daphnia pulex	1.9-2.1 mm	A	S					96	LC <sub>50</sub>	15	Trucco et al., 1983	
Daphnia pulex	≤ 24 h	N	S					48	NOEC	196	Stooff, 1983b	
Daphnia pulex	≤ 24 h	N	S					48	LC <sub>50</sub>	305	Stooff, 1983b	

Table 6.1.1 Single species toxicity data for benzene - continued

Species	Lifestage age/size	A/N	Test system	Purity/ solvent	Testwater	pH	Hardness/ mg CaCO <sub>3</sub> /l	Expo time (h)	Parameter	Results (mg/l)	Reference	Quality
<b>Insects</b>												
<i>Aedes aegypti</i>	3 rd instar	N	S					48	LC <sub>50</sub>	200	Slooff, 1983b	
<i>Culex pipiens</i>	3 rd instar	N	S					48	LC <sub>50</sub>	71	Slooff, 1983b	
<b>Hydrozoans</b>												
<i>Hydra oligactis</i>		N	S					48	LC <sub>50</sub>	34	Slooff, 1983a	
<b>Worms</b>												
<i>Eupobdella octoculata</i>		N	S					48	LC <sub>50</sub>	>320	Slooff, 1983a	
<b>Tubificidae (different species)</b>												
		N	S					48	LC <sub>50</sub>	>320	Slooff, 1983a	
<b>Molluscs</b>												
<i>Lymnaea stagnalis</i>	3-4 wks	N	S					48	NOEC	120	Slooff, 1983a	
<i>Lymnaea stagnalis</i>	3-4 wks	N	S					48	LC <sub>50</sub>	230	Slooff, 1983a	
<b>Echinoderms</b>												
<i>Paracentrotus lividus</i> (N)	Embryo	N	S	Purity >99%				48	EC development	0.078	Pagano, 1988	#2
<i>Paracentrotus lividus</i> (N)	Sperm-offspring	N	S	Purity >99%				48	EC genotoxic	0.078	Pagano, 1988	#2

Table 6.1.1 Single species toxicity data for benzene - continued

Species	Lifestage age/size	A/N	Test system	Purity/ solvent	Testwater	pH	Hardness/ mg CaCO <sub>3</sub> /l	Expo time (h)	Parameter	Results (mg/l)	Reference	Quality
<b>Fish</b>												
<i>Carassus auratus</i>	Juveniles	N	S					96	LC <sub>50</sub>	34.4	Persoone and	
<i>Clupea harengus pallasii</i> (M)	Embryos		S		sal. 2.3%			7d	EC tissue growth delay (reversible)	0.053-0.49	Persoone, 1982	
<i>Clupea harengus pallasii</i> (M)	Spawning females	F			sal. 2.3%			5d	EC respiration	0.053		
								6d	EC embryo development	0.7		
<i>Engraulis mordax</i> (M)	Eggs, 1d-larvae	S			sal. 3%			48	EC development	4-15		
<i>Lepomis macrochirus</i>	Juveniles	N	S					96	LC <sub>50</sub>	22.5	Persoone, 1982	
<i>Morone saxatilis</i>	Juveniles		F		sal. 2.6%			4w	EC growth	5.3	Persoone, 1982	
<i>Oncorhynchus kisutch</i>	Fry	N	S		Freshwater			96	LC <sub>50</sub>	8.6	Persoone, 1982	
<i>Oryzias latipes</i>	4-5 wks	N	S					48	NOEC	126	Slooff, 1983b	
<i>Oryzias latipes</i>	4-5 wks	N	S					48	LC <sub>50</sub>	250	Slooff, 1983b	
<i>Pimephales promelas</i>	Juveniles	N	S		Hard water			96	LOLC	8-9	Persoone, 1982	
<i>Pimephales promelas</i>	Juveniles	N	S		Soft water			96	LC <sub>50</sub>	33.5	Persoone, 1982	
<i>Pimephales promelas</i>	0.58 g	N	F					48	NOEC	54	Slooff, 1983b	
<i>Pimephales promelas</i>	0.58 g	N	F					48	LC <sub>50</sub>	84	Slooff, 1983b	
<i>Poecilia reticulata</i>	Juveniles	N	S					96	LC <sub>50</sub>	36.6	Persoone, 1982	
<i>Poecilia reticulata</i>	3-4 wks	N	S					48	NOEC	265	Slooff, 1983b	
<i>Poecilia reticulata</i>	3-4 wks	N	S					48	LC <sub>50</sub>	420	Slooff, 1983b	
<i>Salmo gairdneri</i>	Juveniles	N	F					96	LC <sub>50</sub>	5.3	Persoone, 1982	
<i>Salmo gairdneri</i>	5-8 wks	N	S					48	NOEC	40	Slooff, 1983b	
<i>Salmo gairdneri</i>	5-8 wks	N	S					48	LC <sub>50</sub>	56	Slooff, 1983b	

Table 6.1.1 Single species toxicity data for benzene - continued

Species	Lifestage age/size	A/N	Test system	Purity/ solvent	Testwater	pH	Hardness/ mg CaCO <sub>3</sub> /l	Expo time (h)	Parameter	Results (mg/l)	Reference	Quality
<b>Amphibians</b>												
<i>Ambystoma americanum</i>	3-4 wks	N	S					48	NOEC	120	Slooff, 1983b	
<i>Ambystoma americanum</i>	3-4 wks	N	S					48	LC <sub>50</sub>	370	Slooff, 1983b	
<i>Xenopus laevis</i>	3-4 wks	N	S					48	NOEC	105	Slooff, 1983b	
<i>Xenopus laevis</i>	3-4 wks	N	S					48	LC <sub>50</sub>	190	Slooff, 1983b	

Table 6.1.2 Single species toxicity data for ethylbenzene

Species	Lifestage age/size	A/N Test system	Purity/ solvent	Testwater	pH	Hardness/ mg CaCO <sub>3</sub> /l	Expo time (h)	Parameter	Results (mg/l)	Reference	Quality
<b>Bacteria</b>											
<i>Pseudomonas putida</i>							16	LOEC cell number	12	Chambon, 1984	
<b>Algae</b>											
<i>Microcystis aeruginosa</i>							8d	LOEC cell number	33	Chambon, 1984	
<i>Scenedesmus quadricauda</i>							8d	LOEC cell number	160	Chambon, 1984	
<i>Selenastrum capricornutum</i>							96	EC <sub>50</sub>	>438	Chambon, 1984	
<i>Skeletonema costatum</i>							96	EC <sub>50</sub>	>438	Chambon, 1984	
<b>Molluscs</b>											
<i>Tapes semidecussata</i> (M)						96	LC <sub>50</sub>	6.8		Chambon, 1984	
<b>Crustaceans</b>											
<i>Artemia</i> (M)	Larvae	S	Purity >97%				24	LC <sub>50</sub>	15.4	Abernethy et al., 1986 #2	
<i>Daphnia magna</i>		S	Purity >97%		6-7		48	LC <sub>50</sub>	2.1	Bobra, 1983 #2	
<i>Daphnia magna</i>	Neonates	S	Purity >97%				48	LC <sub>50</sub>	2.12	Abernethy et al., 1986 #2	
<i>Daphnia magna</i>							24	LC <sub>50</sub>	190	Chambon, 1984	
<i>Daphnia magna</i>							24	LC <sub>50</sub>	77	Chambon, 1984	
<i>Daphnia magna</i>							48	LC <sub>50</sub>	75	Chambon, 1984	
<i>Crangon franciscorum</i> (M)								EC <sub>50</sub>	3.7	Chambon, 1984	
<i>Mysidopsis bahia</i> (M)								EC <sub>50</sub>	87.6	Chambon, 1984	



Table 6.1.2 Single species toxicity data for ethylbenzene - continued

Species	Lifestage age/size	A/N	Test system	Purity/ solvent	Testwater	pH	Hardness/ mg CaCO <sub>3</sub> /l	Expo time (h)	Parameter	Results (mg/l)	Reference	Quality
Mitocra spinipes (M)								24	LC <sub>50</sub>	16	Chambon, 1984	
Palaeomonetes (M) pugio	adult larvae							24	LC <sub>50</sub>	14.5-17.3 10.2	Chambon, 1984	
Fish Carassius auratus								96	LC <sub>50</sub>	94	Chambon, 1984	
Cyprinodon variegatus (M)	Juvenile	N	S	>80%	Seawater (filtered)			96	LC <sub>50</sub> NOEC	280 88	Meitmuller, 1981	
Lepomis macrochirus								96	LC <sub>50</sub>	150	Chambon, 1984	
Morone saxatilis (M)								96	LC <sub>50</sub>	0.43	Chambon, 1984	
Pimephales promelas					soft water hard water hard water			96	LC <sub>50</sub> LC <sub>50</sub> NOEC	48.5 42.3 0.44	Chambon, 1984	
Poecilia reticulata								96	LC <sub>50</sub>	78	Chambon, 1984	
Salmo gairdneri								96	LC <sub>50</sub>	14	Chambon, 1984	

Table 6.1.3 Single species toxicity data for isopropylbenzene

Species	Lifestage age/size	A/N Test system	Purity/ solvent	Testwater	pH	Hardness/ mg CaCO <sub>3</sub> /l	Expo time (h)	Parameter	Results (mg/l)	Reference	Quality
<b>Molluscs</b>											
Mytilus edulis							25d	EC	1	Chambon, 1984	
<b>Crustaceans</b>											
Artemia (M)	Larvae	S	Purity >97%				24	LC <sub>50</sub>	13.7	Abernethy et al., 1986	#2
Daphnia magna	Neonates	S	Purity >97%				48	LC <sub>50</sub>	0.6	Abernethy et al., 1986	#2
Daphnia magna		S	Purity >97%		6-7		48	LC <sub>50</sub>	0.6	Bobra, 1983	#2
Daphnia magna							24	LC <sub>50</sub>	95	Chambon, 1984	

(M): marine organism

- # 1 Well performed experiment with reliable test results; sufficiently documented  
# 2 Experiment with reliable test results; insufficiently documented  
# 3 Experiment with unreliable test results

Table 6.1.4 QSAR estimates of the toxicity of benzene, ethylbenzene and isopropylbenzene to various test species (QSARs from Verhaar and Hermens, 1991). See text for discussion on the reliability.

Species	estimated parameter	Benzene (mg/l)	Ethylbenzene (mg/l)	Isopropylbenzene (mg/l)
<b>Bacteria</b>	NOEC			
<i>Clostridium botulinum</i>	EC <sub>50</sub>	641.2	142.2	61.5
<i>Bacillus subtilis</i>	NOEC	289.2	95.5	51.0
<i>Pseudomonas putida</i>	NOEC	77.8		
<i>Microcystis aeruginosa</i>	NOEC	16.0		
<i>Photobacterium phosphoreum</i>	NOEC	76.5	23.1	11.8
<b>Algae</b>				
<i>Skeletonema costatum</i>	EC <sub>50</sub>	237.7	65.8	32.0
<i>Scenedesmus subspicatus</i>	EC <sub>50</sub>	120.1	24.4	
<i>Selenastrum capricornutum</i>	EC <sub>50</sub>	29.7	4.4	1.5
<b>Protozoa</b>				
<i>Tetrahymena pyriformis</i>	EC <sub>50</sub>	219.2	50.8	22.5
<b>Coelenterates</b>				
<i>Hydra oligactis</i>	NOLC	45.6		
<b>Molluscs</b>				
<i>Lymnaea stagnalis</i>	NOLC	42.6		
<b>Arthropods</b>				
<i>Nitocra spinipes</i>	LC <sub>50</sub>	110.8	26.9	12.2
<i>Daphnia magna</i>	NOEC	8.2	1.1	0.4
<i>Aedes aegypti</i>	LC <sub>50</sub>	139.8		
<i>Aedes aegypti</i>	NOLC	91.6		
<i>Culex pipiens</i>	NOLC	53.6		
<b>Fish</b>				
<i>Alburnus alburnus</i>	LC <sub>50</sub>	135.0	35.0	16.4
<i>Brachydanio rerio</i>	NOEC	14.2	1.8	0.6
<i>Pimephales promelas</i>	LC <sub>50</sub>	41.8	8.7	3.6
<i>Pimephales promelas</i>	NOEC	4.5	0.6	0.2
<i>P.promelas/B.rerio</i>	NOEC	4.3	0.9	0.4
<i>Poecilia reticulata</i>	LC <sub>50</sub>	62.7	12.5	5.1
<b>Amphibia</b>				
<i>Ambystoma mexicanum</i>	NOLC	59.6		
<i>Rana catesbeiana</i>	LC <sub>50</sub>	50.0	10.2	4.2
<i>Rana temporaria</i>	NOLC	54.4		
<i>Xenopus laevis</i>	NOLC	67.9		
<i>Xenopus laevis</i>	LC <sub>50</sub>	15.5		

#### 6.1.4 Quantitative structure activity relations (QSARs)

If sufficient experimental data for a group of structurally related substances are available, quantitative structure activity relations may be established. For "neutral" substances with a narcotic effect, for 25 species QSARs are available where toxicity is a function of the octanol-water partition coefficient (Verhaar and Hermens, 1991). These QSARs may be applied to estimate the toxicity of benzene, ethylbenzene and isopropylbenzene. Table 6.1.4 presents the toxicity data as estimated by the QSAR approach.

For benzene, a comparison of these estimated toxicity data to the experimental data in Tables 6.1.1 to 6.1.3 can be made for 8 species. For 5 species the QSAR

estimate is within a factor 2 of the experimental data and for 3 species the QSAR estimate is 10 to 30 times lower. For ethylbenzene, the QSAR estimate is within a factor 2 for 3 out of 9 test species and for 6 species the QSAR estimate is 6 to 85 times higher than the experimental data. No comparison could be made for isopropylbenzene.

Although the QSAR approach seems to work for benzene, predictions for ethylbenzene are less reliable. If this would be a trend connected to the molecular structure, the approach cannot be applied to supply a reliable estimate for the toxicity of isopropylbenzene.

## **6.2 Toxicity to mammals**

For benzene, ethylbenzene and isopropylbenzene, the relevant routes for exposure of mammals are through oral intake and inhalation. Therefore, the acute toxicity data as presented in Table 6.2.1 are limited mainly to data for oral and inhalation exposure. The acute toxicity of the substances is relatively low. Information on chronic toxicity is presented in the following sections.

### **6.2.1 Benzene**

Benzene caused slight to moderate skin and eye irritation and slight to moderate necrosis (ATSDR, 1989). There are indications that benzene causes fetus-toxicity in animals at 160 mg/m<sup>3</sup> air, but no teratogenic effects or effects on reproduction are found (Slooff, 1988; ATSDR, 1989). A tentative "Tolerable Daily Intake" of 170 µg/kg bodyweight/day was derived by Vermeire et al. (1991).

### **6.2.2 Ethylbenzene**

The toxicity of ethylbenzene was evaluated by Janssen and Van der Heiden (1987). Information on chronic oral toxicity is scant. In one study with rats carried out in 1956, > 400 mg/kg body weight per day (application 5 days a week, during 6 months) caused effects on liver and spleen. No effects were observed at 136 mg/kg body weight (Wolf et al., 1956 in Janssen and Van der Heiden, 1987).

Using a safety factor of 1000 (a high factor was used based on the restricted number and less reliable data) a Tolerable Daily Intake (TDI) of 136 µg/kg body weight/day was derived by Vermeire et al. (1991).

A number of acute inhalation experiments also caused effects on the liver, the lowest concentration causing these effects was 1350 mg/m<sup>3</sup>. No effects were observed at 218 and 430 mg/m<sup>3</sup> (exposure period 6 h a day, 5 days a week). Using a safety factor of 1000, a toxicologically derived acceptable level in air (for continuous exposure) of 77 µg/m<sup>3</sup> was derived (Vermeire et al., 1991).

### **6.2.3 Isopropylbenzene**

Isopropylbenzene is a CNS depressant. In rats exposed to 500 mg/kg daily for 5 months, hyperemia and congestion were noted in the lungs, liver and kidney. Repeated gastric intubation of 154 mg/kg bodyweight did not affect rats, but at 1462 mg/kg, the kidney weight increased (Chambon, 1984).

No new data on the chronic toxicity of isopropylbenzene were found.

Table 6.2.1 Acute toxicity of benzene (Slooff et al., 1986), ethyl benzene and isopropyl benzene (Sax, 1989)

Compound test organism	Route of administration	Expo. time	Criteria	Result (mg/kg)
<b>Benzene</b>				
rat	oral		LD50	5970
rat	oral		LD50	810
rat	oral		LD50	5600
rat	oral		LD50	3000
rat	oral		LD50	3300
rat	oral		LD50	4900
rat	oral		LD50	930
mouse	oral		LD50	4700
rat	inhalation	4 h	LC50	43,700
rat	inhalation	7 h	LC50	31,950
mouse	inhalation	2 h	LC50	60,000
mouse	inhalation	8 h	LC50	15,000
<b>ethylbenzene</b>				
rat	oral		LD50	3,500
mouse	intraperitoneal		LD50	2,272
guinea pig, adult	inhalation		LCLo	10,000
human	inhalation	8 h	TCLo Eye, Nerv., Pulm.	100
rat	inhalation	7 h	TCLo Repr.	97
rat	inhalation	7 h	TCLo Ter.	985
rat	inhalation	7 h	TCLo Ter.	96
rat	inhalation	4 h	LCLo	4,000
mouse	inhalation	2 h	LCLo	50
rabbit	skin		LD50	17,800
<b>isopropylbenzene</b>				
rat	oral		LD50	1,400
rat	inhalation	4 h	LC50	8,000
mouse	inhalation	2 h	LC50	24,700
rabbit	skin		LD50	12,300

\* mg/l Repr. = reproduction effects  
 \*\* mg/m<sup>3</sup> Ter. = teratogene effects  
 \*\*\* g/m<sup>3</sup> Eye = eye effects  
 Nerv. = nervous-system effects  
 Pulm. = pulmonary system effects

## 6.3 Carcinogenicity, mutagenicity and teratogenicity

The Dutch National Institute of Public Health and Environmental Protection (RIVM) has evaluated the effects of benzene, and ethylbenzene for deriving toxicological criteria (Vermeire et al., 1991). The most important results are summarized below.

### 6.3.1 Benzene

Benzene is considered to have genotoxic properties, although there is no clear evidence of a direct reaction with DNA. In cells inheritable chromosomal aberrations causing neoplastic transformation were shown. Benzene was found to be carcinogenic in rodents after oral and inhalation exposure.

Chronic exposure of man to benzene in air at concentrations  $\geq 325$  mg/m<sup>3</sup> are known to induce leukaemia, mostly initiated with non-neoplastic effects on the blood forming system.

Chronic exposure of man to concentrations over 100 mg/m<sup>3</sup> air causes aberrations of the hematopoietical system. Effects are anaemia, leucopenia and thrombocytopenia, and in severe cases aplastic anaemia.

Because of its carcinogenic activity in man it is concluded that the risk assessment for benzene should be based on a non-threshold approach. By means of linearly extrapolation of reliable data it is calculated that life-time exposure through inhalation of  $12 \mu\text{g}/\text{m}^3$  or oral administration of ca.  $120 \mu\text{g}$  per day (100% absorption) causes a risk of  $1 : 10^4$ . As the working mechanism for the occurrence of anaemia in man is unknown, it is assumed that similar concentrations of benzene administered via oral and inhalation exposure causes comparable effects. According to the Dutch Public Health organization the linear model leads to a strong overestimation of the real risk. They conclude that mean concentrations below  $1.200 \mu\text{g}/\text{m}^3$  (ca.  $12.000 \mu\text{g}/\text{day}$  intake) do not exceed the risk of  $1 : 10^4$ .

### **6.3.2 Ethylbenzene and isopropylbenzene**

The carcinogenity, mutagenity and teratogenity of ethylbenzene and isopropylbenzene cannot be evaluated due to a lack of data.

## **7 ENVIRONMENTAL IMPACT ASSESSMENT**

### **7.1 Comparison of exposure and effects**

After release into the environment benzene, ethylbenzene and isopropylbenzene will mainly partition to the atmosphere because of their high volatilization rate. Few data are available on sorption and bioaccumulation. The chemical degradation rate in water will be low as a result of the stability of these structures. However, benzene and isopropylbenzene are classified as readily biodegradable. Therefore, biodegradation in the aquatic environment is expected to occur. In the atmosphere the substances will degrade rapidly by indirect photolysis with radicals.

Observed environmental concentrations for benzene, ethylbenzene and isopropylbenzene were reported to vary from 3 µg/l to below detection limits in river water and seawater. Concentrations in the sediment were unknown.

The relatively low log  $K_{ow}$ -values (2.1, 3.2 and 3.8) do not indicate strong bioaccumulative properties. Moreover, experimental data for benzene showed that bioconcentration in fish was lower than predicted on the basis of the lipophilicity (log  $K_{ow}$ ). Yet low levels of ethylbenzene were reported in fish (4 µg/kg in the liver) and in crab.

The toxicity of these substances varies roughly between 1 and 200 mg/l. For benzene, this is well below the water solubility, but for ethylbenzene (150 mg/l) and isopropylbenzene (50 mg/l), the water solubility is in the same range.

The lowest reported concentration without effect in an acute test is 40 mg/l for trout. However, effects were observed at lower concentrations in other tests; therefore available No Observed Effect Concentrations cannot be taken as a basis for deriving water quality criteria. An effective concentration for the most sensitive species was 0.08 mg/l in an embryo development test with an echinoderm. The margin of safety between these toxicity data and the above mentioned common environmental levels is at least a factor 25.

### **7.2 Water quality standard**

For the estimation of a 'safe' level for the environment, an extrapolation factor can be applied to the available toxicity data which accounts for the different sensitivity of other, non-tested species in the ecosystem. Various approaches have been proposed to extrapolate from single species toxicity data to a concentration where the aquatic ecosystem is not affected (OECD, 1991). This  $NOEC_{ecosystem}$  may also be called maximal tolerable concentration or MTC.

The 'modified US-EPA method' can be applied even if only one  $LC_{50}$ -value is available. The method applies an assessment factor of 10 to account for each additional level of uncertainty: (1) from one or several chronic values to the  $NOEC_{ecosystem}$ , (2) from acute  $LC_{50}$ -values for at least algae, Daphnia and fish to a chronic  $NOEC$ -value, and (3) from only one acute  $LC_{50}$ -value to a chronic value. Thus, the assessment factors are 10, 100 and 1000, respectively.

Another approach is to use the variability in the sensitivity among the various test species as a means to calculate a concentration that is expected to be safe for most (e.g. 95%) of the species in the aquatic ecosystem. In other words, a concentration is calculated that is hazardous for only a small number (5%) of species. Two calculation methods are available, which differ in their basic assumptions regarding the shape of the distribution curve for the species sensitivity. One method assumes a log-logistic distribution (Aldenberg and Slob, 1991), whereas the other assumes a log-normal distribution (Wagner and Løkke, 1991).

The results of the two main approaches are presented in Table 7.2.1. The extrapolation method based on the difference in species sensitivity are very sensitive for extreme values. The inclusion of a NOEC of 600 mg/l for an algae results in an unrealistically low MTC for benzene. Therefore the value calculated by the EPA method (lowest NOEC/10; in this case the NOEC is a LOEC/2) is preferred as MTC: 0.005 mg/l.

Table 7.2.1 Maximum tolerable concentrations for benzene

Extrapolation method	MTC (mg/l)	Remarks
Modified EPA	0.004	lowest NOEC / 10
Aldenberg and Slob, 1991	0.0008	{ 95% protection level, (50% confidence,) 5 test species
Wagner and Løkke, 1991	0.0004	

It is assumed that the toxicity of benzene, ethylbenzene and isopropylbenzene is related to their log  $K_{ow}$ . Therefore the toxicity of benzene is expected to be 27 times lower than the toxicity of ethylbenzene and 47 times lower than isopropylbenzene. However, the intraspecies variability for benzene is higher than this difference among substances. Therefore it is proposed to calculate a MTC for benzene and assume that the value is valid for the sum of the three substances.

Another safety factor should account for the combined effects of many other xenobiotic chemicals present in the environment. These benzene substitutes and many other 'industrial' chemicals are neutral substances that show narcosis-type toxicity. The effects of these substances are supposed to be additive (Hermens et al., 1984). The order of magnitude for this safety factor is unknown at the present state of the art. One could suggest a safety factor of 10 when not many other environmental contaminants are expected, but when the environmental load is high, a safety factor of 1000 could be more appropriate.

The magnitude of this safety factor is not only a matter of science but political considerations are involved as well. Therefore no safety factor is applied to the MTC to derive a water quality standard for the sum of benzene, ethylbenzene and isopropylbenzene of 5 µg/l (rounded off). It should be realized that the proposed water quality standard does not take into account that other industrial pollutants may be present at the same time.

This proposed level is detectable by various analytical techniques as described in chapter 2.

For benzene the U.S. has proposed a Maximum Concentration Limit for humans of 5 µg/l. Furthermore a health advisory is proposed at 0.1 µg/l concerning the risk of cancer. For ethylbenzene the U.S. has proposed Maximum Concentration Limit of 700 µg/l for lifetime exposure.

Canadian guidelines (1987) for benzene and ethylbenzene are 5 µg/l and 2.4 µg/l (Joint Research Centre, 1989). Standards of the USA and Canada are comparable with the proposed water quality standards (5 µg/l).

However, the standard proposed by the USA for ethylbenzene of 700 µg/l is more than two orders of a magnitude higher.



In the previous report to EC (Persoone and Vanhaecke, 1982) limit concentrations of 200 and 100 µg/l were proposed to protect freshwater biota and the marine environment against benzene, respectively. In view of the recent approaches in hazard assessment, however, these limits can no longer be maintained.

When the proposed water quality standard is compared to the environmental levels commonly found in the riverwater of 0.2 to 3 µg/l for benzene and 0.7 to 2.3 for ethylbenzene (see Chapter 3.1.3) the levels come close to the proposed water quality standard.

### 7.3 Human exposure

Exposure of man to substances in the aquatic environment may occur through oral intake from (drinking) water and from fish, shellfish or crayfish. Other possible exposure routes are dermal contact and inhalation. The route of dermal contact is relevant if a substance is lipophilic, whereas inhalation is important for highly volatile substances. For the sake of simplicity, intake by inhalation and dermal contact are considered to be negligible in relation to the oral uptake for benzene, ethylbenzene and isopropylbenzene. The human exposure analysis is therefore exclusively based on oral intake.

For the calculations the total concentration of benzene, ethylbenzene and isopropylbenzene in the water is assumed to be at the level of the proposed water quality standard: 5 µg/l. Assuming a daily water consumption of 2 litre, the oral intake of total benzenes from the water will be 10 µg/day.

The concentration in fish is estimated from the bioconcentration factor ( $C_{\text{fish}}/C_{\text{water}} \leq 10$ ) to be  $\leq 100$  µg/kg fish. If furthermore a (relatively high) figure for the daily consumption of fish is assumed to be 100 g, the oral intake will be  $\leq 10$  µg/day.

The total oral intake of total benzenes is estimated at 20 µg/day or for a 60 kg person 0.33 µg/kg body weight/day. Comparison with the "tentative acceptable daily intake" for benzene of 6.5 µg/kg body weight/day and for ethylbenzene of 136 µg/kg body weight/day (see chapter 6.3) the margin of safety is rather high. Therefore the risk of human exposure to benzene, ethylbenzene and isopropylbenzene present in aquatic systems in a concentration of 5 µg/l is expected to be negligible.

## REFERENCES

- Abernethy S., A.M. Bobra, W.Y. Shiu, P.G. Mackay and D. Wells, 1986.  
Acute lethal toxicity of hydrocarbons and chlorinated hydrocarbons to two planktonic crustaceans: the key role of organism-water partitioning.  
*Aquatic Toxicology* 8, 163-174.
- Aldenberg, T. W. Slob, 1991  
Confidence limits for hazardous concentrations based on logistically distributed NOEC toxicity data.  
National Institute of Public Health and Environmental Protection (RIVM), no. 71902002.
- Arvin E., B.K. Jensen and A.T. Gundersen, 1989.  
Substrate interactions during aerobic biodegradation of benzene.  
*Appl. Environ. Microbiol.*, 55(12), 3221-3225.
- ATSDR, 1989.  
Toxicological profile for benzene.  
Atlanta, US, Agency for Toxic Substance and Disease Registry, Centers for Disease Control, Public Health. Service, Department of Health & Human Services, ATSDR/TP-88/03.
- Bobra A.M., W.Y. Shiu and D. Mackay, 1983.  
A predictive correlation for the acute toxicity of hydrocarbons and chlorinated hydrocarbons to the water flea (*Daphnia magna*).  
*Chemosphere* 12 (9/10), 1121-1129.
- Bouwer E.J., 1989.  
Biotransformation of aromatics in strip-pit pond.  
*J. Environ. Engineering*, 115(4).
- Bouwer E.J. and P.L. McCarty, 1983.  
Transformations of halogenated organic compounds under denitrification conditions.  
*Appl. Environ. Microbiol.* 45, 1295-1299.
- Bouwer E.J. and P.L. McCarty, 1985.  
Utilization rates of trace halogenated organic compounds in acetate grown biofilms.  
*Biotechnol. Bioeng.* 27, 1564-1571, from Keuning and Janssen, 1987.
- Bouwer E.J. 1985.  
Secondary utilization of trace halogenated organic compounds in biofilms.  
*Environ. Progr.* 4, 43-46, from Keuning and Janssen, 1987.
- Casserly D.M., E.M. Davis, T.D. Downs and R.K. Guthrie, 1983.  
Sorption of organisms by *Selenastrum capricornatum*.  
*Wat. Res.*, 17, 11, 1591-1594.
- Chambon P., 1984.  
Evaluation of the impact of ethylbenzene and cumene on the aquatic environment

De Bruin, J., 1985.

Model classification scheme for information pertaining to dangerous substances which could be included in list I of the council directive 76/464 and the establishment of a system to collect, store and use such information (ref. nr. XI/A/3).

De Bruijn J., 1991.

Hydrophobicity, biokinetics and toxicity of environmental pollutants: A Structure Activity approach, 168 pp.

Delfino J.J. and C.J. Miles, 1985.

Aerobic and anaerobic degradation of organic contaminants in Florida groundwater.

Proceedings/Soil and Crop Science Society of Florida 44, 9-14, from Keuning and Janssen, 1987.

ECETOC, 1988.

Concentrations of industrial organic chemicals measured in the environment: The influence of physico-chemical properties, tonnage and use pattern.

Technical report 29.

EURECO, 1990.

Technical datasheets on substances, candidates for List I, Directive 76/464/EEC.

EPA, 1981.

An exposure and risk assessment for benzene, EPA-440/4-85-006.

EPA, 1984.

EPA Method Study 25, method 602, purgeable aromatics, EPA-600/4-84-042.

Geyer R.A. (ed.), 1980.

Marine Environmental Pollution, 1 (hydrocarbons), Elsevier Oceanography Series, 27A, Amsterdam.

Gesellschaft Deutscher Chemiker, 1988.

Benzol. BUA-Stoffbericht 24. Weinheim, New York.

From EURECO, 1990.

Gibson D.T., B. Gschwendt, W.K. Yeh and V.M. Kobal, 1973.

Initial reactions in the oxidation of ethylbenzene by *Pseudomonas putida*.

Biochemistry 12, 1520-1528, from Keuning and Janssen, 1987.

Gibson D.T., 1972.

The microbial oxidation of aromatic hydrocarbons.

CRC Crit. Rev. Microbiol. 1, 199-223, from Keuning and Janssen, 1987.

Gibson D.T. and V. Subramanian, 1984.

Microbial degradation of aromatic hydrocarbons, p. 188.

In: Microbial degradation of organic compounds Ed. D.T. Gibson. Dekker Inc., New York, from Keuning and Janssen, 1987.

- Heitmuller P.T., T.A. Hollister and P.R. Parrish, 1981.  
Acute Toxicity of 54 Industrial Chemicals to Sheepshead Minnows (*Cyprinodon variegatus*).  
Bull. Environm. Contam. Toxicol. 27, 596-604.
- Hellmann H., 1987.  
Modellversuche zur Verflüchtigung organischer Spurenstoffe in Oberflächen-  
gewässern, Fresenius Z. Anal. Chem., 328, 475-479.
- Hermens J., P. Leeuwangh and A. Mush, 1984  
Quantitative structure-activity relationships and mixture toxicity studies of chloro-  
and alkylanilines at acute lethal toxicity level to the guppy *Poecilia reticulata*.  
Ecotox. Environ. Saf., 9, 17-25.
- Holland H.L., I.M. Carter, P.C. Chenchiah, S.H. Kahn, B. Munoz, R.W.  
Ninnis and D. Richards, 1985.  
Fungal hydroxylation of ethylbenzene and derivatives.  
Tetrahedron Lett. 26, 6409-6412, from Keuning and Janssen, 1987.
- Hutchins S.R., G.W. Sewell, D.A. Kovacs and G.A. Smith, 1991.  
Biodegradation of aromatic hydrocarbons by aquifer micro-organisms under  
denitrifying conditions.  
Environ. Sci. Technol. 25, 68-76.
- Joint Research centre, 1989.  
Scientific assessment of EC standards for drinking water quality, environment and  
quality of life.  
Commission of the European Communities Cat. No. CD-NA-12427-EN-C, 188-  
197.
- IDET-CEGOS. C.B. Belin, 1983.  
Aspect Technico-économiques des mesures de réduction de la pollution des eaux  
par les rejets de certain dérivés chlorés du benzène. Final report submitted to the  
Commision of the EC. Contract No. 84/B/6600/11/023/11/S; Boulogne-  
Billancourt.  
From EURECO, 1990.
- Ince N., Y. Inel, 1989.  
Volatilization of organic chemicals from water.  
Water, Air, and Soil Pollution, 47, 71-79.
- Janssen P. en C. van der Heijden, 1987.  
Korte samenvatting en evaluatie van de toxicologie van ethylbenzeen  
(Geaccordeerd door de ACT-adviesgroep toxicologie op 24 maart 1987). RIVM,  
Bilthoven.
- Karlson U. and W.T. Frankenberger Jr., 1989.  
Microbial degradation of benzene and toluene in groundwater.  
Bull. Environ. Contam. Toxicol. 43, 505-510.
- Keuning S. en D.B. Janssen, 1987.  
Microbiologische afbraak van zwarte en prioritaire stoffen voor het milieubeleid.  
Biochemisch Laboratorium, Rijksuniversiteit Groningen, The Netherlands.

- Kincannon D.F., A. Weinert, R. Padorr and E.L. Stover, 1983.  
Predicting treatability of multiple organic priority pollutant wastewaters from single-pollutant treatability studies.  
Proc. Ind. Waste Conf. 37, 641-650, from Keuning and Janssen, 1987.
- Mackay D., P.J. Leikonen, 1975.  
Rate of evaporation of low-solubility contaminants from water bodies to atmosphere.  
Environ. Sci. Technol., 9, 1178-1180.
- Major D.W., C.I. Mayfield and J.F. Barker, 1988.  
Biotransformation of benzene by denitrification in aquifer sand.  
Ground Water 26(1), 8-14.
- Melcer H., H. Monteith, M. Newbigging, S.G. Nutt and J.P. Stephenson, 1989.  
Dynamic modelling of trace contaminant removal in a municipal sewage treatment plant.  
Wat. Sci. Tech. 21, 351-362.
- Meyer J.S., M.D. Marcus and H.L. Bergman, 1984.  
Inhibitory interactions of aromatic organics during microbial degradation.  
Environ. Tox. Chem. 3, 583-587.
- Mill T., D.G. Hendry, H. Richardson, 1980.  
Free-radicals oxidants in natural water: photooxidation of Cumene and Pyridine.  
Science, 207 (4433), 886-887.
- MITI, Ministry of International Trade and Industry, Japan, 1985.  
List existing chemical substances tested on biodegradation by micro-organisms or bioaccumulation in fish body by MITI, Annex.
- OECD, 1991.  
Guidance document for aquatic effects assessment.  
Draft report 60 pp.
- Oldenhuis R., L. Kuyk, A. Lumrners, D.B. Janssen and B. Witholt, 1989.  
Degradation of chlorinated and non-chlorinated aromatic solvents in soil suspensions by pure bacterial cultures.  
Appl. Microbiol. Biotechnol. 30, 211-217.
- Pagano G., M. Cipollaro, G. Corsale, A. Esposito, G.G. Giordano, E. Ragucci, M. Trieff, 1988.  
Comparative toxicities of benzene, chlorobenzene and dichlorobenzenes to sea urchin embryos and sperm.  
Bull. Environ. Contam. Toxicol. 40, 481-488.
- Persoone G., P. Vanhaecke, 1982.  
Evaluation of the impact of benzene, chloroform and carbon tetrachloride on the aquatic environment. EC-report no. XI/627/82.
- Rogerson A., W.Y. Shiu, G.L. Huang, D. MacKay and J. Berger, 1983.  
Determination and interpretation of hydrocarbon toxicity to ciliate protozoa.  
Aquatic Toxicology 3, 215-228.

- Schink B., 1985.  
Degradation of unsaturated hydrocarbons by methanogenic enrichment c  
FEMS Microbiology Ecology 31, 69-77.
- Sittig M., 1981.  
Hanbook of toxic and hazardous chemicals.  
ISBN 08155-0841-7.
- Slooff W., 1983a.  
Benthic macroinvertebrates and water quality assessment: some toxicological considerations.  
Aquatic Toxicology 4, 73-82.
- Slooff W., J.H. Canton and J.L.M. Hermens, 1983b.  
Comparison of the susceptibility tests of 22 freshwater species to 15 chemical compounds. I (sub)acute toxicity tests.  
Aquatic Toxicology 4, 113-128.
- Slooff W. et al., 1986.  
Basisdocument benzeen.  
RIVM report no. 840760002.
- Slooff W., (ed.), 1988.  
Basisdocument benzeen,  
Bilthoven, Rijksinstituut voor Volksgezondheid en Milieuhygiëne, Publikatiereeks Milieubeheer, serie basisdocumenten.
- Smith M.R. and C. Ratledge, 1989.  
Catabolism of alkylbenzenes by *Pseudomonas* sp. NCIB 10643.  
Appl. Microbiol. Biotechnol. 32, 68-75.
- Thursby G.B., R.L. Steele and M.E. Kane, 1985.  
Effect of organic chemicals on growth and reproduction in the marine red alga *Champia parvula*.  
Environ. Toxicol. and Chem. 4, 797-805.
- Trucco R.G., F.R. Engelhardt B. Stacey, 1983.  
Toxicity, accumulation and clearance of aromatic hydrocarbons in *Daphnia pulex*  
Environ. Poll. (Series A) 31, 191-202.
- US-EPA, 1984.  
Health Assessment Document for Chlorinated Benzenes.  
Washington, Office of Health and Environmental Assessment, Environment Protection Agency, review draft EPA-600/8-84-015A.
- Utkin I.B., M.M. Yakimov, L.N. Matveeva, E.I. Kozlyak, I.S. Royozhin, Z.G. Solomon and A.M. Bezborodov, 1991.  
Degradation of styrene and ethylbenzene by *Pseudomonas* species Y2.  
FEMS Microbiology Letters 77, 237-242.

- Verhaar H.J.M and J.L.M. Hermens, 1991.  
Quantitative structure-activity relationships of environmental pollutants: an outline for application within hazard assessment of new and existing chemicals. Research Institute of Toxicology, University of Utrecht, EEC/DBXI, Contract B6614/90/3040.
- Vermeire T.G., M.E. van Apeldoorn, J.C. de Fouw, en P.J.C.M. Janssen, 1991.  
Voorstel voor de humaan-toxicologische onderbouwing van C-(toetsings)waarden. Rijksinstituut voor Volksgezondheid en Milieuhygiëne, report no. 725201005, 170 pp.
- Vogel T.M. and D. Grbic-Galic, 1986.  
Incorporation of oxygen from water into toluene and benzene during anaerobic fermentive transformation.  
Appl. Environ. Microbiol. 52, 200-202, from Keuning and Janssen, 1987.
- Wagner C. and H. Løkke, 1991.  
Estimation of ecotoxicological protection levels from NOEC toxicity data.  
Wat. Res. 25, 10, 1237-1242.
- Ware G.W. (ed.), 1980.  
Reviews of Environmental Contamination and Toxicology, Volume 106, Springer-Verlag, New York.
- Water Research Centre, 1987.  
The importance of diffuse pollution sources and their consequences to the environment of the Community. Final report submitted to the Commission of the EC.  
From EURECO, 1990.
- WHO, 1983.  
L'arsenic. critères d'hygiène de l'environnement 18. Genève.  
From EURECO, 1990.
- Wilson B.H., G.B. Smith and J.F. Rees, 1986.  
Biotransformation of selected alkylbenzenes and halogenated aliphatic hydrocarbons in methanogenic aquifer material: a microcosm study.  
Environ. Sci. Technol. 20, 997-1002, from Keuning and Janssen, 1987.
- Wolf, M.A., V.K. Rowe, D.D. McCollister, R.L. Hollingsworth en F. Oyen, 1956.  
Toxicological studies of certain alkylated benzenes and benzene.  
Arch. Ind. Health, 14, 387-398.
- Wong P.I.S., Y.K. Chau, J.S. Rhamey and M. Docker, 1984.  
Relationship between water solubility of chlorobenzenes and their effects on a freshwater green alga.  
Chemosphere 13, 991-996.

## APPENDIX 1.

The calculation method for the volatilization standard half-life according to De Bruin (1985).

In the calculation method some assumptions have been made:

- water flow rate = 1 m/s
- air velocity = 3 m/s
- water depth = 1 m
- temperature = 10 °C

The calculation consists of 6 different steps that will be described below:

### 1 Henry Constant (H):

$$H = p/s$$

$p$  = vapour pressure (atm.)  
 $s$  = water solubility (mol/m<sup>3</sup>)  
 $H$  = Henry Constant (atm·m<sup>3</sup>/mol)

### 2 dimensionless Henry constant (H'):

$$H' = H/RT$$
$$RT = 0.024$$

### 3 Liquid-Phase Exchange Constant (k(l)):

if  $M < 65$ :

$$k(l) = 20\sqrt{44/M}$$

if  $M > 65$ :

$$k(l) = 41.93\sqrt{32/M}$$

$M$  = molecular weight

### 4 Gas-Phase Exchange Constant (k(g)):

$$k(g) = 4550\sqrt{18/M}$$

### 5 Overall liquid-phase mass transfer coefficient (K(L)):

$$K(L) = \{H' \cdot k(g) \cdot k(l)\} / \{H' \cdot k(g) + k(l)\}$$

### 6 Half-life ( $t_{1/2}$ ):

$$t_{1/2} = 69.3/K(L)$$

$t_{1/2}$  = half-life (hours)



---

# Eastern Europe and the USSR

---

## THE CHALLENGE OF FREEDOM


---

GILES MERRITT

The sparks of unrest that leapt from Berlin in November 1989 to Moscow's Red Square in August 1991 are firing an explosion of political and economic change. Out of the ashes of Communism is emerging the shape of a vast new European market-place stretching from the Atlantic to the Pacific.

In his fascinating account of Europe's fast-changing East-West relationships, Giles Merritt argues that a massive rescue operation must be mounted to ensure the success of these changes. The upheaval of Communism's collapse is 'The challenge of freedom'.

Written with the cooperation and support of the European Commission, this book sets out to identify the key policy areas where a new partnership is being forged between the countries of Eastern and Western Europe. It offers a privileged insight into the current thinking of European



Community officials, politicians and industrial leaders, and analyses the factors that will determine whether the emerging market economies of Eastern Europe can truly be absorbed into a single European economy.

Immensely readable and often disturbing, this important book contains much up-to-date and hitherto unpublished information on such major East-West problem areas as energy, environmental control, immigration, trade relations, agriculture and investment. It also examines the arguments surrounding a 'Marshall Plan' for Eastern Europe that would emulate the famous US aid programme that helped relaunch the economies of Western Europe in the aftermath of World War II.

For anyone concerned about the future of Eastern Europe and the USSR, whether from a political, social or economic standpoint, this book is essential reading.

# Bulletin of the European Communities



The *Bulletin of the European Communities*, which is issued 10 times a year (monthly, except for the January/February and July/August double issues), is an official reference publication covering all spheres of Community activity.

It is compact, easy to consult (with an index and copious references to the Official Journal and to previous issues), logically structured (to reflect the main fields of Community policy) and wholly reliable. The Bulletin is an essential reference tool, describing the passage of Community legislation through all its stages from presentation of a proposal by the Commission to final enactment by the Council.

Thanks to its topical commentaries on the month's major events, it provides the student of European integration and other interested readers with up-to-date and accurate information about the most recent developments in Community policy — the creation of a single market, economic and social integration, the Community's role in international affairs, etc.

Supplements to the Bulletin are published from time to time, containing important background material on significant issues of the day. Recent Supplements have covered German unification, the Commission's programme for 1992 and European industrial policy for the 1990s.

The Bulletin and its Supplements are produced by the Secretariat-General of the Commission, 200 rue de la Loi, B-1049 Brussels, in the nine official languages of the Community, and can be ordered from the Community sales agents.



# INFO92

## The Community database focusing on the objectives and the social dimension of the single market

As a practical guide to the single market, INFO92 contains vital information for all those determined to be ready for 1992.

INFO92 is really a simple market scoreboard, recording the state of play on the stage-by-stage progress of Commission proposals up to their adoption by the Council, summarizing each notable development and placing it in context, and keeping track of the transposition of directives into Member States' national legislation.

Using INFO92 is simplicity itself. It can be consulted on-screen by means of a wide range of everyday equipment connected to specialized data-relay networks. Fast transmission, the virtually instant updating facility (several times a day, if necessary) and dialogue procedures requiring no prior training make INFO92 ideal for the general public as well as for business circles and the professions.

The system offers easy access to information thanks to the choice of menus available and to the logical presentation modelled on the structure of the *White Paper*, the *Social Charter* and the decision-making process within the institutions.

Enquiries may also be made to the Commission Offices in the Member States or – for small businesses – the Euro-Info Centres now open in all regions of the Community.

---

**Eurobases Helpdesk**

{ Tel. : (32-2) 295 00 03  
Fax : (32-2) 296 06 24

# **DIRECTORY**

## **OF COMMUNITY LEGISLATION IN FORCE and other acts of the Community institutions**

The Community's legal system is of direct concern to the individual citizen as much as to the Member States themselves.

Both lawyers and non-lawyers, then, need to be familiar not just with national law, but also with Community legislation, which is implemented, applied or interpreted by national law and in some cases takes precedence over it.

To make Community legislation more accessible to the public, the Commission of the European Communities publishes a Directory, updated twice a year, covering:


- binding instruments of secondary legislation arising out of the Treaties establishing the three Communities (regulations, decisions, directives, etc.);
- other legislation (internal agreements, etc.);
- agreements between the Communities and non-member countries.

Each entry in the Directory gives the number and title of the instrument, together with a reference to the Official Journal in which it is to be found. Any amending instruments are also indicated, with the appropriate references in each case.

The legislation is classified by subject matter. Instruments classifiable in more than one subject area appear under each of the headings concerned.

The Directory proper (Vol. I) is accompanied by two indexes (Vol. II), one chronological by document number and the other alphabetical by keyword.

The Directory is available in the nine official languages of the Community.



1 064 pp. – ECU 83  
ISBN 92-77-77093-7 (Volume I)  
ISBN 92-77-77094-5 (Volume II)  
ISBN 92-77-77095-3 (Volume I and II)  
FX-86-91-001-EN-C  
FX-86-91-002-EN-C

# EUROPEAN ECONOMY


*European Economy* appears four times a year, in March, May, July and November. It contains important reports and communications from the Commission to the Council and to Parliament on the economic situation and developments, as well as on the borrowing and lending activities of the Community. In addition, *European Economy* presents reports and studies on problems concerning economic policy.

Two supplements accompany the main periodical:

- Series A – ‘Economic trends’ appears monthly except in August and describes with the aid of tables and graphs the most recent trends of industrial production, consumer prices, unemployment, the balance of trade, exchange rates, and other indicators. This supplement also presents the Commission staff’s macroeconomic forecasts and Commission communications to the Council on economic policy.
- Series B – ‘Business and consumer survey results’ gives the main results of opinion surveys of industrial chief executives (orders, stocks, production outlook, etc.) and of consumers (economic and financial situation and outlook, etc.) in the Community, and other business cycle indicators. It also appears monthly, with the exception of August.

Unless otherwise indicated, the texts are published under the responsibility of the Directorate-General for Economic and Financial Affairs of the Commission of the European Communities, 200 rue de la Loi, B-1049 Brussels, to which enquiries other than those related to sales and subscriptions should be addressed.

Subscription terms are shown on the back cover and the addresses of the sales offices are shown on the third page of the cover.



# Success in business


**depends on the decisions you make ...  
which depend on the information you receive**

Make sure that your decisions are based on information that is accurate and complete!

In a period of rapid adjustment, with national economies merging into a single European economy under the impetus of 1992, reliable information on the performance of specialized industry sectors is essential to suppliers, customers, bankers and policymakers.

Small and medium-sized enterprises, in particular, need easy access to information.

The market must be defined, measured and recorded. Information is needed on production capacities, bottlenecks, future developments, etc.




**Panorama of EC industry 1991-1992**  
**Current situation and outlook for 180 sectors**  
**of manufacturing and service industries**  
**in the European Community**

1 400 pp. \* ECU 110 \* ISBN 92-826-3103-6 \* CO-60-90-321-EN-C

# SOCIAL EUROPE

*Social Europe*, published by the Commission of the European Communities, Directorate-General for Employment, Industrial Relations and Social Affairs (DG V), Coordination and Information Policy Unit, deals with current social affairs in Europe.

The basic review appears three times a year. In addition, a number of supplements/files are published annually, each dealing in depth with a given subject.



# ENERGY

## A CHALLENGE FOR EUROPE AND THE WORLD

---


Since it first appeared in 1985 **Energy in Europe** has become recognized as an invaluable source of information on both the policy-making and the operational aspects of European Community energy policy. Subscribers include leaders of energy-consuming and energy-producing industries and other decision-makers in the private and public sectors, as well as major consultancies and research institutes in and outside the Community.

In the present situation within the Community, itself at the eve of the single market, and *vis-à-vis* the huge energy problems, as well as the potential, of our neighbours in Central and Eastern Europe and in the Commonwealth of Independent States, the energy sector is of the greatest strategic importance. An understanding of it is indispensable in many areas of economic activity. It also constitutes a crucial factor within a debate of truly global importance, namely the protection of the environment, including the global warming issue.

**Energy in Europe** continues to keep its readers abreast of the ongoing situation as regards overall policy, markets, energy planning, and the constant quest for cleaner and more efficient energy technology.

Market trends and perspectives are covered in **two regular issues** each year, and also in a **Short-term energy outlook** appearing in the first half of the year and an **Annual energy review** at the end of the year which includes the world energy situation by region including EC Member States, the short-term energy outlook for the Community, and a review of trends in main indicators over 10 years. Further **Special Issues** are also produced in connection with major developments or events, including international conferences on or relevant to the energy sector.

**Energy in Europe** appears in English but each issue also contains translations into French, German or Spanish of articles from the preceding issue.





Also available:

---

**Treaty on European Union**

253 pp. \* ECU 9 \* ISBN 92-824-0959-7 \* RX-73-92-796-EN-C — 1992

---

**Conradh ar an Aontas Eorpach**

253 pp. \* ECU 9 \* ISBN 92-824-0964-3 \* RX-73-92-796-GA-C — 1992

---

**National implementing measures to give effect to the White Paper of the Commission on the completion of the internal market  
*Situation at 31 October 1992***

448 pp. \* ECU 47 \* ISBN 92-826-4973-3 \* CM-76-92-875-EN-C — 1992

---

**Reports of Commission Decisions relating to competition – 1989/1990**

247 pp. \* ECU 12 \* ISBN 92-826-3868-5 \* CV-73-92-772-EN-C — 1992

---

**Harmonization of company law in the European Community –  
Measures adopted and proposed – *Situation as at 1 March 1992***

518 pp. \* ECU 54 \* ISBN 92-826-4314-X \* C1-74-92-831-EN-C — 1992

---

**Green Paper on the development of the single market for postal services (Communication from the Commission) – COM(91) 476 final**

371 pp. \* ISBN 92-77-45007-X \* CB-CO-92-263-EN-C — 1992

---

**The European Community and human rights**

61 pp. \* ECU 8 \* ISBN 92-826-5083-9 \* CM-76-92-407-EN-C — 1993

---

**The opening-up of public procurement**

46 pp. \* ECU 6 \* ISBN 92-826-5130-4 \* CO-77-92-085-EN-C — 1993

---

**Farm take-over and farm entrance within the EEC**

148 pp. \* ECU 15 \* ISBN 92-826-3667-4 \* CM-73-91-376-EN-C — 1992

---

**Copyright and information limits to the protection of literary and pseudo-literary works in the Member States of the EC**

262 pp. \* ECU 27 \* ISBN 92-826-3666-6 \* CM-75-92-049-EN-C — 1992

---

**Social security for persons moving within the Community –  
*Social Europe 3/92***

154 pp. \* ECU 19 \* ISSN 0255-0776 \* CB-AA-92-003-EN-C — 1992

---

---

**Urban social development – *Social Europe – Supplement 1/92***

125 pp. \* ECU 9 \* ISBN 92-826-4013-2 \* CE-NC-92-001-EN-C — 1992

---

**Enterprise and people aspects in the information technology sector to the year 2000 – *Social Europe – Supplement 2/92***

111 pp. \* ECU 9 \* ISBN 92-826-4537-1 \* CE-NC-92-002-EN-C — 1992

---

**Towards a Europe of solidarity: housing – *Social Europe – Supplement 3/92***

142 pp. \* ECU 9 \* ISBN 92-826-4567-3 \* CE-NC-92-003-EN-C — 1992

---

**The regulation of working conditions in the Member States of the European Community – Volume 1 – *Social Europe – Supplement 4/92***

114 pp. \* ECU 9 \* ISBN 92-826-4996-2 \* CE-NC-92-004-EN-C — 1992

---

**XXVth General Report on the Activities of the European Communities – 1992**

520 pp. \* ECU 20 \* ISBN 92-826-5340-4 \* CM-76-92-681-EN-C — 1993

---

**EUR 13914 – European cooperation in the field of scientific and technical research,  
COST secretariat with the assistance of J. L. Roland**

93 pp. \* ECU 16.50 \* ISBN 92-826-4371-9 \* CG-NA-13914-EN-C — 1992

---

**EUR 14006 – Thesaurus guide – *Second edition, prepared by EUROBrokers***

1033 pp. \* ECU 78 \* ISBN 92-826-4956-3 \* CD-NA-14006-EN-C — 1993

---

**EUR 14197 – Evaluation of economic effects: relevance and impacts of EC programmes promoting industrial R&D with special emphasis on small and medium-sized enterprises (pilot methodological study)  
K. HORNSCHILD, F. MEYER-KRAHMER**

129 pp. \* ECU 13.50 \* ISBN 92-826-3817-0 \* CD-NA-14197-EN-C — 1992

---

**EUR 14198 – Evaluation of the impact of European Community research programmes upon the competitiveness of European industry – concepts and approaches**

J.S. METCALFE, L. GEORGHIOU, P. CUNNINGHAM, H.M. CAMERON

44 pp. \* ECU 6.00 \* ISBN 92-826-3818-9 \* CD-NA-14198-EN-C — 1992

---

**EUR 14326 – The European market for value analysis**

139 pp. \* ECU 15 \* ISBN 92-826-4130-9 \* CD-NA-14326-EN-C — 1992

---

---

## **Employment in Europe – 1992**

206 pp. \* ECU 11.25 \* ISBN 92-826-4559-2 \* CE-75-92-106-EN-C — 1992

---

## **Focus on the East – *Energy in Europe***

157 pp. \* ECU 19 \* ISSN 1017-6705 \* CS-BI-92-001-4H-C — 1992

---

## **A view to the future – *Energy in Europe***

176 pp. \* ECU 19 \* ISBN 92-826-3665-8 \* CS-75-92-841-EN-C — 1992

---

## **The finances of Europe, Daniel STRASSER**

439 pp. \* ECU 18.50 \* ISBN 92-826-2306-8 \* CM-60-90-280-EN-C — 1992

---

## **XXIst Report on Competition Policy – 1991**

446 pp. \* ECU 29 \* ISBN 92-826-3850-2 \* CM-73-92-247-EN-C — 1992

---

## **Europe in figures – *Third edition***

256 pp. \* ECU 16.50 \* ISBN 92-826-3371-1 \* CA-70-91-895-EN-C — 1992

---

## **Inventory of taxes levied in the Member States of the European Communities – *14th edition***

726 pp. \* ECU 80 \* ISBN 92-826-0417-9 \* CM-59-90-855-EN-C — 1992

---

## **A common market for services – *Current status 1 january 1993***

*Banking • Insurance • Transactions in securities • Transport services • New technologies and services • Capital movements • Free movement of labour and the professions*

(Internal market – Volume 1)

184 pp. \* ECU 19 \* ISBN 92-826-5257-2 \* CO-10-93-001-EN-C — 1993

---

## **The elimination of frontier controls – *Current status 1 january 1993***

*Control of goods • Control of individuals • Value-added tax • Excise duties*

(Internal market – Volume 2)

128 pp. \* ECU 19 \* ISBN 92-826-5268-8 \* CO-10-93-002-EN-C — 1993

---

## **Conditions for business cooperation – *Current status 1 january 1993***

*Company law • Intellectual property • Company taxation – Public procurement*

**Internal market for energy**

(Internal market – Volume 3)

104 pp. \* ECU 19 \* ISBN 92-826-5277-7 \* CO-10-93-003-EN-C — 1993

---

---

**Community social policy – Labour market • Employment and pay • Improved living and working conditions • Free movement of workers • Social protection • Freedom of association and collective bargaining • Information, consultation and participation of employees • Equal treatment for men and women • Vocational training • Health and safety at work • Rights and protection of children and adolescents • The elderly • The disabled**  
(Completing the internal market – Volume 6)

308 pp. \* ECU 34 \* ISBN 92-826-3609-7 \* CO-62-91-006-EN-C — 1992

---

**The current situation, evolution and future prospects  
for agriculture in Yugoslavia**

128 pp. \* ECU 12 \* ISBN 92-826-3485-X \* CM-72-91-899-EN-C — 1992

---

**A practical guide to cross-border cooperation**

112 pp. \* ECU 10 \* ISBN 92-826-3143-5 \* CT-70-91-992-EN-C — 1992

---

**1992: a pivotal year** (address by Jacques Delors, President of the Commission, to the European Parliament) – **From the Single Act to Maastricht and beyond: the means to match our ambitions – The Commission's programme for 1992** – *Supplement 1/92 – Bull. EC*

45 pp. \* ECU 5 \* ISBN 92-826-3841-3 \* CM-NF-92-001-EN-C — 1992

---

**Research after Maastricht: an assessment, a strategy –**  
*Supplement 2/92 – Bull. EC*

49 pp. \* ECU 5 \* ISBN 92-826-4307-7 \* CM-NF-92-002-EN-C — 1992

---

**Europe and the challenge of enlargement –** *Supplement 3/92 – Bull. EC*

24 pp. \* ECU 5 \* ISBN 92-826-4524-X \* CM-NF-92-003-EN-C — 1992

---

**The creation of the internal market in insurance, Bill POOL**

126 pp. \* ECU 10.50 \* ISBN 92-826-0246-X \* CB-58-90-336-EN-C — 1992

---

**European Economy – No 44 – One market, one money –**  
**An evaluation of the potential benefits and costs of forming an  
economic and monetary union**

351 pp. \* ECU 18 \* ISSN 0379-0991 \* CB-AR-90-044-EN-C — 1992

---

**European Economy – The economics of EMU – Background studies  
for European Economy No 44 'One market, one money'**  
*Special edition No 1 – 1991*

248 pp. \* ECU 18 \* ISBN 92-826-1996-6 \* CM-60-90-208-EN-C — 1992

---

---

**European Economy – No 51 – The climate challenge – Economic aspects of the Community's strategy for limiting CO<sub>2</sub> emissions**

253 pp. \* ECU 23.50 \* ISSN 0379-0991 \* CM-AR-92-051-EN-C — 1992

---

**Practical guide to legal aspects of industrial subcontracting within the European Community – Volume I – The subcontract**

118 pp. \* ECU 11.25 \* ISBN 92-825-9593-5 \* CB-27-89-001-EN-C — 1992

---

**Practical guide to legal aspects of industrial subcontracting in the European Community – Volume II – The legal framework of subcontracting in the twelve Member States**

70 pp. \* ECU 9 \* ISBN 92-826-4651-3 \* CB-27-89-002-EN-C — 1992

---

**The rules governing medicinal products in the European Community – Volume I – The rules governing medicinal products for human use in the European Community – Revised edition / September 1991**

228 pp. \* ECU 20 \* ISBN 92-826-3166-4 \* CO-71-91-631-EN-C — 1992

---

**The rules governing medicinal products in the European Community – Volume II – Notice to applicants for marketing authorizations for medicinal products for human use in the Member States of the European Community**

186 pp. \* ECU 16.50 \* ISBN 92-825-9503-X \* CB-55-89-293-EN-C — 1992

---

**The rules governing medicinal products in the European Community – Volume III – Addendum No 2 – May 1992 – Guidelines on the quality, safety and efficacy of medicinal products for human use**

206 pp. \* ECU 23 \* ISBN 92-826-4550-9 \* CO-75-92-558-EN-C — 1992

---

**The rules governing medicinal products in the European Community – Volume IV – Good manufacturing practice for medicinal products**

177 pp. \* ECU 19.50 \* ISBN 92-826-3180-X \* CO-71-91-760-EN-C — 1992

---

**The rules governing medicinal products in the European Community – Volume V – Veterinary medicinal products**

158 pp. \* ECU 14.25 \* ISBN 92-825-9643-5 \* CB-55-89-972-EN-C — 1992

---

---

**The rules governing medicinal products in the European Community – Volume VI – Establishment by the European Community of maximum residue limits (MRLs) for residues of veterinary medicinal products in foodstuffs of animal origin**

127 pp. \* ECU 13.50 \* ISBN 92-826-3173-7 \* CO-71-91-768-EN-C — 1992

---

**Credit Institutions – Community measures adopted or proposed – Situation as at August 1992**

356 pp. \* ECU 37 \* ISBN 92-826-4842-7 \* CI-76-92-293-EN-C — 1992

---

**New information technology in education – France**

110 pp. \* ECU 13 \* ISBN 92-826-4771-4 \* CY-03-92-007-EN-C — 1992

---

**New information technology in education – Germany**

124 pp. \* ECU 13 \* ISBN 92-826-4683-1 \* CY-03-92-004-EN-C — 1992

---

**New information technology in education – Denmark**

138 pp. \* ECU 14 \* ISBN 92-826-4681-5 \* CY-03-92-003-EN-C — 1992

---

**New information technology in education – The Netherlands**

108 pp. \* ECU 12 \* ISBN 92-826-4779-X \* CY-03-92-011-EN-C — 1992

---

**New information technology in education – Luxembourg**

122 pp. \* ECU 13 \* ISBN 92-826-4777-3 \* CY-03-92-010-EN-C — 1992

---

**Removal of tax obstacles to the cross-frontier activities of companies – Supplement 4/91 – Bull. EC**  
67 pp. \* ECU 4.25 \* ISBN 92-826-3025-0 \* CM-NF-91-004-EN-C — 1992

**Agriculture in Europe**

97 pp. \* ECU 7 \* ISBN 92-826-0476-4 \* CM-60-90-418-EN-C — 1992

**Telecommunications in Europe**, Herbert UNGERER with the collaboration of Nicholas P. COSTELLO  
*Revised edition, 1990*

257 pp. \* ECU 10 \* ISBN 92-826-1640-1 \* CM-59-90-346-EN-C — 1992

**European Economy – No 35 – The economics of 1992**

222 pp. \* ECU 16 \* ISSN 0379-0991 \* CB-AR-88-035-EN-C — 1992

**European Economy – No 40 – Horizontal mergers and competition policy in the European Community**

98 pp. \* ECU 16 \* ISSN 0379-0991 \* CB-AR-89-040-EN-C — 1992

**European Economy – No 45 – Stabilization, liberalization and devolution – Assessment of the economic situation and reform process in the Soviet Union**

191 pp. \* ECU 18 \* ISSN 0379-0991 \* CB-AR-90-045-EN-C — 1992

**European Economy – No 50 – Annual Economic Report 1991-92 – Strengthening growth and improving convergence**

285 pp. \* ECU 20 \* ISSN 0379-0991 \* CM-AR-91-050-EN-C — 1992

**European Economy – Social Europe – The impact of the internal market by industrial sector: the challenge for the Member States – Special edition – 1990**

340 pp. \* ECU 18 \* ISBN 92-826-1818-8 \* CM-59-90-887-EN-C — 1992

**Social Europe 3/91 – Equal opportunities for women and men**

202 pp. \* ECU 18 \* ISSN 0255-0776 \* CE-AA-91-003-EN-C — 1992

---

---

**Guide to the reform of the Community's structural Funds**

104 pp. \* ECU 11.25 \* ISBN 92-826-0029-7 \* CB-56-89-223-EN-C — 1992

**The rights of working women in the European Community, Eve C. LANDAU**

244 pp. \* ECU 5.25 \* ISBN 92-825-5341-8 \* CB-43-85-741-EN-C — 1992

**European Economy – No 36 – Creation of a European financial area – Liberalization of capital movements and financial integration in the Community**

212 pp. \* ECU 16 \* ISSN 0379-0991 \* CB-AR-88-036-EN-C — 1992

**1992: the European social dimension, Patrick VENTURINI**

119 pp. \* ECU 9.75 \* ISBN 92-825-8703-7 \* CB-PP-88-B05-EN-C — 1992

**Research on the 'cost of non-Europe' – Basic findings**

**Volume 1 – Basic studies: Executive summaries**

578 pp. \* ECU 53.25 \* ISBN 92-825-8605-7 \* CB-PP-88-B14-EN-C — 1992

**Volume 2 – Studies on the economics of integration**

652 pp. \* ECU 57 \* ISBN 92-825-8616-2 \* CB-PP-88-C14-EN-C — 1992

**Volume 3 – The completion of the internal market:**

**A survey of European industry's perception of the likely effects**

309 pp. \* ECU 25.50 \* ISBN 92-825-8610-3 \* CB-PP-88-D14-EN-C — 1992

**Volume 4 – The 'cost of non-Europe':**

**Border-related controls and administrative formalities –**

**An illustration in the road haulage sector**

280 pp. \* ECU 22.50 \* ISBN 92-825-8618-9 \* CB-PP-88-E14-EN-C — 1992

**Volume 5 (Parts A + B)**

**The 'cost of non-Europe' in public-sector procurement**

Part A: 552 pp. \* ISBN 92-825-8646-4 \* CB-P1-88-F14-EN-C — 1992

Part B: 278 pp. \* ISBN 92-825-8647-2 \* CB-P2-88-F14-EN-C — 1992

Parts A + B: ECU 120 \* ISBN 92-825-8648-0

**Volume 6 – Technical barriers in the EC: An illustration by six industries**

**The 'cost of non-Europe': Some case studies on technical barriers**

242 pp. \* ECU 21 \* ISBN 92-825-8649-9 \* CB-PP-88-G14-EN-C — 1992

**Volume 7 – The 'cost of non-Europe': Obstacles to transborder business activity**

154 pp. \* ECU 12.75 \* ISBN 92-825-8638-3 \* CB-PP-88-H14-EN-C — 1992

**Volume 8 – The 'cost of non-Europe' for business services**

140 pp. \* ECU 13.50 \* ISBN 92-825-8637-5 \* CB-PP-88-I14-EN-C — 1992

**Volume 9 – The 'cost of non-Europe' in financial services**

494 pp. \* ECU 120 \* ISBN 92-825-8636-7 \* CB-PP-88-J14-EN-C — 1992

**Volume 10 – The benefits of completing the internal market for telecommunication services equipment in the Community**

197 pp. \* ECU 17.25 \* ISBN 92-825-8650-2 \* CB-PP-88-K14-EN-C — 1992

**Volume 11 – The EC 92 automobile sector**

350 pp. \* ECU 27.75 \* ISBN 92-825-8619-7 \* CB-PP-88-L14-EN-C — 1992

**Volume 12 (Parts A + B)**

**The 'cost of non-Europe' in the foodstuffs industry**

Part A: 424 pp. \* ISBN 92-825-8642-1 \* CB-P1-88-M14-EN-C — 1992

Part B: 328 pp. \* ISBN 92-825-8643-X \* CB-P2-88-M14-EN-C — 1992

Parts A + B: ECU 120 \* ISBN 92-825-8644-8

**Volume 13 – «Le coût de la non-Europe» des produits de construction**

168 pp. \* ECU 14.25 \* ISBN 92-825-8631-6 \* CB-PP-88-N14-FR-C — 1992

**Volume 14 – 'The cost of non-Europe' in the textile-clothing industry**

256 pp. \* ECU 21.75 \* ISBN 92-825-8641-3 \* CB-PP-88-O14-EN-C — 1992

**Volume 15 – 'The cost of non-Europe' in the pharmaceutical industry**

182 pp. \* ECU 13.50 \* ISBN 92-825-8632-4 \* CB-PP-88-P14-EN-C — 1992

**Volume 16 – The internal markets of North America – Fragmentation and integration in the USA and Canada**

176 pp. \* ECU 13.50 \* ISBN 92-825-8630-8 \* CB-PP-88-Q14-EN-C — 1992

*Special price for the complete series: ECU 360*

# Commission of the European Communities

## OFFICES

---

### **EIRE — IRELAND**

#### ***Dublin***

Commission of the European Communities  
Office in Ireland

Jean Monnet Centre  
39 Molesworth Street  
Dublin 2  
Tel. (353-1) 71 22 44  
Fax (353-1) 71 26 57  
Telex (0500) 93827 EUCO EI

### **UNITED KINGDOM**

#### ***London***

Commission of the European Communities  
Office in the United Kingdom

Jean Monnet House  
8 Storey's Gate  
London SW1P 3AT  
Tel. (44-71) 973 19 92  
Fax (44-71) 973 19 00/19 10  
Telex (051) 23208 EURUK G

#### ***Belfast***

Commission of the European Communities  
Office in Northern Ireland

Windsor House  
9/15 Bedford Street  
Belfast BT2 7EG  
Tel. (44-232) 24 07 08  
Fax (44-232) 24 82 41  
Telex (051) 74117 CECBEL G

#### ***Cardiff***

Commission of the European Communities  
Office in Wales

4 Cathedral Road  
Cardiff CF1 9SG  
Tel. (44-222) 37 16 31  
Fax (44-222) 39 54 89  
Telex (051) 497727 EUROPA G

#### ***Edinburgh***

Commission of the European Communities  
Office in Scotland

9 Alva Street  
Edinburgh EH2 4PH  
Tel. (44-31) 225 20 58  
Fax (44-31) 226 41 05  
Telex (051) 727420 EUEDING

### **UNITED STATES OF AMERICA**

#### ***Washington***

Commission of the European Communities  
External Delegation

2100 M Street, NW (Suite 707)  
Washington DC 20037  
Tel. (1-202) 862 95 00  
Fax (1-202) 429 17 66  
Telex (023) 64215 EURCOM NW

#### ***New York***

Commission of the European Communities  
External Delegation

3 Dag Hammarskjöld Plaza  
305 East 47th Street  
New York NY 10017  
Tel. (1-212) 371 38 04  
Fax (1-212) 758 27 18  
Telex (023) 012396 EURCOM NY

### **JAPAN**

#### ***Tokyo***

Commission of the European Communities  
External Delegation

Europa House  
9-15 Sanbancho  
Chiyoda-Ku  
Tokyo 102  
Tel. (81-3) 239 04 41  
Fax (81-3) 239 93 37  
Telex (072) 28567 COMEUTOK J



**BELGIQUE/BELGIË****Bruxelles/Brussel**

Commission des Communautés  
européennes  
Commissie van de Europese  
Gemeenschappen  
Bureau en Belgique/Bureau in België  
Rue Archimèdestraat 73  
B-1040 Bruxelles/Brussel  
Tél.: (32-2) 295 38 44  
Fax: (32-2) 295 01 66  
Télex: 26657 COMINF B

**DANMARK****København**

Kommissionen for De Europæiske  
Fællesskaber  
Kontor i Danmark  
Højbrohus  
Østergade 61  
Postbox 144  
DK-1004 København K  
Tlf.: (45-33) 14 41 40  
Telefax: (45-33) 11 12 03/14 13 92  
Telex: (055) 16402 COMEUR DK

**FRANCE****Paris**

Commission des Communautés  
européennes  
Bureau de représentation en France  
288, boulevard Saint-Germain  
F-75007 Paris  
Tél.: (33-1) 40 63 38 00  
Fax: (33-1) 45 56 94 17/18/19  
Télex: (042) 202271F

**Marseille**

Commission des Communautés  
européennes  
Bureau à Marseille  
2, rue Henri-Barbusse  
F-13241 Marseille Cedex 01  
Tél.: (33) 91 91 46 00  
Fax: (33) 91 90 98 07  
Télex: (042) 402538 EURMA

**LUXEMBOURG**

Commission des Communautés  
européennes  
Bureau au Luxembourg  
Bâtiment Jean Monnet  
Rue Alcide De Gasperi  
L-2920 Luxembourg  
Tél.: (352) 43 01  
Fax: (352) 43 01 44 33  
Télex: 3423/3446/3476 COMEUR LU

**SUISSE/SCHWEIZ/SVIZZERA****Genève**

Commission des Communautés  
européennes  
Bureau de presse et d'information  
Kommission der Europäischen  
Gemeinschaften  
Presse- und Informationsbüro

Commissione delle Comunità europee  
Ufficio stampa e informazione  
Case postale 195  
37-39, rue de Vermont  
CH-1211 Genève 20 CIC  
Tél.: (41-22) 734 97 50  
Fax: (41-22) 734 22 36  
Télex: (045) 414 165 ECOM CH

**BR DEUTSCHLAND****Bonn**

Kommission der Europäischen  
Gemeinschaften  
Vertretung in der Bundesrepublik  
Deutschland  
Zitelmannstraße 22  
D-53113 Bonn 1  
Tél.: (49-228) 53 00 90  
Fax: (49-228) 530 09 50/12  
Telex: (041) 886648 EUROP D

**Berlin**

Kommission der Europäischen  
Gemeinschaften  
Vertretung in der Bundesrepublik  
Deutschland  
Außenstelle Berlin  
Kurfürstendamm 102  
D-W-10711 Berlin 31  
Tél.: (49-30) 896 09 30  
Fax: (49-30) 892 20 59  
Telex: (041) 184015 EUROP D

**München**

Kommission der Europäischen  
Gemeinschaften  
Vertretung in der Bundesrepublik  
Deutschland  
Vertretung in München  
Erhardtstraße 27  
D-80331 München 2  
Tél.: (49-89) 202 10 11  
Fax: (49-89) 202 10 15  
Telex: (041) 5218135

**ESPAÑA****Madrid**

Comisión de las Comunidades  
Europeas  
Oficina en España  
Calle de Serrano, 41, 5ª planta  
E-28001 Madrid  
Tél.: (34-1) 435 17 00  
Telecopia: (34-1) 576 03 87/577 29 23  
Télex: (052) 46818 OIPE E

**Barcelona**

Comisión de las Comunidades  
Europeas  
Oficina en España  
Avenida Diagonal, 407 bis, 18ª planta  
E-08008 Barcelona  
Tél.: (34-3) 415 81 77  
Telecopia: (34-3) 415 63 11  
Télex: (052) 97524 BDC E

**VENEZUELA****Caracas**

Comisión de las Comunidades  
Europeas  
Delegación  
Calle Orinoco, Las Mercedes  
Apartado 67 076,  
Las Américas 1061A  
Caracas  
Tél.: (58-2) 91 51 33  
Telecopia: (58-2) 91 88 76  
Télex: (031) 27298 COMEU VC

**ΕΛΛΑΔΑ****Αθήνα**

Επιτροπή των Ευρωπαϊκών  
Κοινοτήτων  
Γραφείο στην Ελλάδα  
Βασιλίσσης Σοφίας 2  
ΤΘ 30284  
GR-10674 Αθήνα  
Τηλ.: (30-1) 724 39 82/83/84  
Τέλεφαξ: (30-1) 724 46 20  
Τελεξ: (0601) 219324 ECAT GR

**ITALIA****Roma**

Commissione delle Comunità europee  
Ufficio in Italia  
Via Poli, 29  
I-00187 Roma  
Tél.: (39-6) 699 11 60  
Telecopia: (39-6) 679 16 58 679 36 52  
Telex: (043) 610184 EUROMA I

**Milano**

Commissione delle Comunità europee  
Ufficio a Milano  
Corso Magenta, 59  
I-20123 Milano  
Tél.: (39-2) 48 01 25 05  
Telecopia: (39-2) 481 85 43  
Telex: (043) 316200 EURMIL I

**NEDERLAND****Den Haag**

Commissie van de Europese  
Gemeenschappen  
Bureau in Nederland  
Korte Vijverberg 5  
2513 AB Den Haag  
Nederland  
Tél.: (31-70) 346 93 26  
Telecopie: (31-70) 364 66 19  
Telex: (044) 31094 EURCO NL

**PORTUGAL****Lisboa**

Comissão das Comunidades  
Europeas  
Gabinete em Portugal  
Centro Europeu Jean Monnet  
Largo Jean Monnet, 1-10.  
P-1200 Lisboa  
Tél.: (351-1) 350 98 00  
Telecopiador: (351-1) 350 98 01/2/3  
Telex: (0404) 18810 COMEUR P

## European Communities — Commission

### Updating of data concerning the impact on the aquatic environment of certain dangerous substances, second part — Part I

#### Document

Luxembourg: Office for Official Publications of the European Communities

1993 — XII, 41 pp. — 21.0 x 29.7 cm

Part I: ISBN 92-826-5894-5

Parts I-IV: ISBN 92-826-5893-7

Price (excluding VAT) in Luxembourg: Part I: ECU 7  
Parts I-IV: ECU 34

Venta y suscripciones • Salg og abonnement • Verkauf und Abonnement • Πωλήσεις και συνδρομές  
Sales and subscriptions • Vente et abonnements • Vendita e abbonamenti  
Verkoop en abonnementen • Venda e assinaturas

<b>BELGIQUE / BELGIË</b>  Monteur belge / Belgisch Staatsblad Rue de Louvain 42 / Leuvenseweg 42 B-1000 Bruxelles / B-1000 Brussel Tél. (02) 512 00 26 Fax (02) 511 01 84  Autres distributeurs / Overige verkooptenten Librairie européenne/ Europese boekhandel Rue de la Loi 244/Vetstraat 244 B-1040 Bruxelles / B-1040 Brussel Tél. (02) 231 04 35 Fax (02) 735 08 60  Jean De Lanoy Avenue du Roi 202 / Koningslaan 202 B-1060 Bruxelles / B-1060 Brussel Tél. (02) 538 51 89 Télex 63220 UNBOOK B Fax (02) 538 08 41  Document delivery: Credoc Rue de la Montagne 34 / Bergstraat 34 Bte 11 / Bus 11 B-1000 Bruxelles / B-1000 Brussel Tél. (02) 511 89 41 Fax (02) 513 31 95	<b>FRANCE</b>  Journal officiel Service des publications des Communautés européennes 26, rue Desaix F-75727 Paris Cedex 15 Tél. (1) 40 58 75 00 Fax (1) 40 58 77 00  <b>IRELAND</b>  Government Supplies Agency 4-5 Harcourt Road Dublin 2 Tel. (1) 66 13 111 Fax (1) 47 80 645  <b>ITALIA</b>  Licosa SpA Via Duca di Calabria 1/1 Casella postale 552 I-50125 Firenze Tel. (055) 84 54 15 Fax 84 12 57 Telex 570466 LICOSA I  <b>GRAND-DUCHÉ DE LUXEMBOURG</b>  Messagerie du livre 5, rue Raiffaesen L-2411 Luxembourg Tél. 40 10 20 Fax 40 10 20 01  <b>NEDERLAND</b>  SDU Overheidsinformatie Externe Fondsen Postbus 20014 2500 EA 's-Gravenhage Tel. (070) 37 99 911 Fax (070) 34 75 778  <b>PORTUGAL</b>  Imprensa Nacional Casa da Moeda, EP Rua D. Francisco Manuel de Melo, 5 P-1062 Lisboa Codex Tel. (01) 68 34 14  Distribuidora de Livros Bertrand, Lda Grupo Bertrand, SA Rua das Terras dos Vales, 4-A Apartado 37 P-2700 Amadora Codex Tel. (01) 49 59 050 Telex 15798 BERDIS Fax 49 60 255  <b>UNITED KINGDOM</b>  HMSO Books (Agency section) HMSO Publications Centre 51 Nine Elms Lane London SW8 5DR Tel. (071) 873 9090 Fax 873 9463 Telex 29 71 138  <b>ÖSTERREICH</b>  Manzsche Verlags- und Universitätsbuchhandlung Kohnmarkt 16 A-1014 Wien Tel. (0222) 531 61-133 Telex 112 500 BOX A Fax (0222) 531 61-181  <b>SUOMI/FINLAND</b>  Akateeminen Kirjakauppa Keskuskatu 1 PO Box 126 SF-00101 Helsinki Tel. (0) 121 41 Fax (0) 121 44 41	<b>NORGE</b>  Narvesen Info Center Bertrand Narvesens vei 2 PO Box 6125 Etterstad N-0602 Oslo 6 Tel. (22) 57 33 00 Telex 79666 NIC N Fax (22) 98 19 01  <b>SVERIGE</b>  STJ AB Traktorvägen 13 S-22100 Lund Tel. (046) 18 00 00 Fax (046) 18 01 25 30 79 47  <b>SCHWEIZ / SUISSE / SVIZZERA</b>  OSEC Stampfenbachstraße 65 CH-8035 Zürich Tel. (01) 385 54 49 Fax (01) 385 54 11  <b>ČESKÁ REPUBLIKA</b>  NIS ČR Havelská 22 130 00 Praha 3 Tel. (2) 235 84 46 Fax (2) 235 97 88  <b>MAGYARORSZAG</b>  Euro-Info-Service Club Sziget Margitsziget 1138 Budapest Tel./Fax 1 111 60 61 1 111 62 16  <b>POLSKA</b>  Business Foundation ul. Krucza 38/42 00-512 Warszawa Tel. (22) 21 99 93, 628-28-62 International Fax&Phone (0-39) 12-00-77  <b>ROMANIA</b>  Euromedia 65, Strada Dionisie Lupu 70184 Bucuresti Tel./Fax 0 12 96 46  <b>BALGARİJA</b>  Europress Klassica BK Ltd 66, bd Vitosha 1463 Sofia Tel./Fax 2 52 74 75  <b>RUSSIA</b>  GCEC 9,60-Istiya Oktyabrya Avenue 117112 Moscow Tel./Fax (095) 135 52 27  <b>CYPRUS</b>  Cyprus Chamber of Commerce and Industry Chamber Building 38 Graves Digenis Ave 3 Delgiorgis Street PO Box 1455 Nicosia Tel. (2) 49500/462312 Fax (2) 458630  <b>MALTA</b>  Miller distributors Ltd Scots House, M.A. Vassalli street PO Box 272 Valletta Tel. 24 73 01 Fax 23 49 14	<b>TÜRKİYE</b>  Pres Gazete Kitap Dergi Pazarlama Degitim Ticaret ve Sanayi AŖ Naribahçe Sokak N. 15 İstanbul-Çağaloğlu Tel. (1) 520 92 96 - 528 55 66 Fax 520 64 57 Telex 23822 DSVO-TR  <b>ISRAEL</b>  ROY International PO Box 13056 41 Mishmar Hayarden Street Tel Aviv 61130 Tel. 3 496 108 Fax 3 544 60 39  <b>UNITED STATES OF AMERICA / CANADA</b>  UNIPUB 4611-F Assembly Drive Lanham, MD 20706-4391 Tel. Toll Free (800) 274 4888 Fax (301) 459 0056  <b>CANADA</b>  Subscriptions only Uniquement abonnements Renouf Publishing Co. Ltd 1294 Algoma Road Ottawa, Ontario K1B 3W8 Tel. (613) 741 43 33 Fax (613) 741 54 39 Telex 0534783  <b>AUSTRALIA</b>  Hunter Publications 58A Gipps Street Collingwood Victoria 3066 Tel. (3) 417 5361 Fax (3) 419 7154  <b>JAPAN</b>  Kinokuniya Company Ltd 17-7 Shinjuku 3-Chome Shinjuku-Ku Tokyo 160-91 Tel. (03) 3439-0121  Journal Department PO Box 55 Chitose Tokyo 156 Tel. (03) 3439-0124  <b>SOUTH-EAST ASIA</b>  Legal Library Services Ltd STK Agency Robinson Road PO Box 1817 Singapore 9036  <b>SOUTH AFRICA</b>  Safto 5th Floor, Export House Cnr Maude & West Streets Sandton 2146 Tel. (011) 883-3737 Fax (011) 883-6569  <b>AUTRES PAYS OTHER COUNTRIES ANDERE LANDE</b>  Office des publications officielles des Communautés européennes 2, rue Mercier L-2985 Luxembourg Tél. 499 28-1 Telex PUBOF LU 1324 b Fax 48 85 73/48 68 17
--	--	---	---

---

Price (excluding VAT) in Luxembourg: Part I: ECU 7  
Parts I-IV: ECU 34

ISBN 92-826-5894-5

 OFFICE FOR OFFICIAL PUBLICATIONS  
OF THE EUROPEAN COMMUNITIES

L-2985 Luxembourg

  
9 789282 658949 >